

5. Silicates-Volume 1

#56

9/28/73

SILICATES VOL I #56

Q 26

VOLUME 1

GRAS MONOGRAPH SERIES

SILICATES

prepared for
THE FOOD AND DRUG ADMINISTRATION
DEPARTMENT OF HEALTH, EDUCATION
AND WELFARE

SEPTEMBER 28, 1973

prepared by
Tracor Jitco, Inc.

Table of Contents

Summary

CHEMICAL INFORMATION

	Page
I. Nomenclature	1
II. Empirical Formula	2
III. Structural Formula	2
IV. Molecular Formula	2
V. Specifications	8
VI. Description	14
VII. Analytical Methods	14
VIII. Occurrence	17
IX. Stability in Containers	22

BIOLOGICAL DATA

I. Acute Toxicity	23
Animal Studies	23
Humans	26
II. Short-Term Studies	28
Rats	28
Guinea Pigs	35
Dogs	37
Humans	39
III. Long-Term Studies	41
Rabbits	41
Humans	43
IV. Special Studies	47
Carcinogenicity	47
Teratogenic Effects	52
Effects on Action of Bacilli	53

BIOCHEMICAL ASPECTS

I. Breakdown	55
II. Absorption and Distribution	55
Rats	55
Rabbits	60
Humans	61
III. Metabolism and Excretion	65
Rats	66
Guinea Pigs	68
Rabbits	72
Dogs	72
Humans	75
IV. Effects on Enzymes and Other Biochemical Parameters	83
Renal Mechanisms and Lesions	84
Serum Protein and Biosynthesis Alterations	87
Blood Lipids	89
Enzyme Changes	90
Effects on Blood Cells and Clotting	91
V. Drug Interaction	92
VI. Mechanism of Toxicity and the Factors Affecting It	96
VII. Consumer Exposure Information	105

SUMMARY

Description

Silica and the silicates from sources that are likely to be candidates for human intake occur in a number of different forms -- 30 to 40 as a rough estimation. They exhibit a range of solubilities and of crystalline or noncrystalline forms. Generally speaking most are assumed to be somewhat soluble from the surface area in the juices of the gastrointestinal tract.

Occurrence

In the natural environment, silicon occurs in practically all forms of plant, animal, and sea life (262). It is in all natural waters: mineral water, hot springs, city water, well-water and sea water (006). Also, it has been found in the atmosphere in the vicinity of factories manufacturing silicon and its alloys (147). As a food additive, up to 2% may be incorporated in a given food (018,019,020 and 021). Average daily intakes have been estimated as 481 mg for calcium silicate, 340 mg for silica serogel, 2278 mg for sodium aluminum silicate, and 2.66 mg for sodium calcium aluminosilicate hydrate (375).

Acute Toxicity

Lethal Dose (LD) values (from various forms of silica) of from 9 mg/kg BW (194) to 150 mg/kg BW (209) for mice by the intravenous route have been reported. Intraperitoneally, from 50 mg/kg BW to 250 mg/kg for silica sol (209) and 4250 mg/kg

for talc administered to mice are recorded. LD values for rats range from 35.2 mg/kg (084) to 100 mg/kg (209) i.v., intratracheally 30 and 50 mg/kg (397), and intraperitoneally 100 mg/kg for these animals (397).

For guinea pigs i.p. 100 (397) and 120 to 240 mg/kg (209), and i.v. 100 mg/kg (209) are recorded. Rabbits given i.v. injections succumbed to doses from 35 to 141 mg/kg (209), cats from 15 to 193 mg/kg (348), and dogs intragastrically 25-50 cc of colloidal silicic acid (476) and i.v. 20 mg/kg of particulate silica (260). In humans, fatal cases from talc aspiration (351), and talc embolism from i.v. injections by drug addicts (043,542) have been reported. Short term studies on rats have shown positive weight-gain effects (061) with no toxic effects from diatomaceous earth of large particle size. Weight-gain effects were variable for rats fed various forms of commercial silicic acid preparations (273). Precipitated amorphous silica (of particle size 19-25 mu) administered intratracheally to rats (084) resulted in lung lesions. Silicates; aluminum, sodium or magnesium, along with silicon dioxide gave only mild clinical symptoms of polydipsia and polyuria when fed to rats, with no drug-related lesions noted (372). Dogs, however, treated similarly (372) showed gross

Short Term
Studies

and microscopic renal lesions from the sodium and magnesium silicates, but none with aluminum silicate or silicon dioxide. Guinea pigs administered a variety of talc samples i.p. showed a fibrosis reaction similar to the foreign body reaction (441). Children exposed to low atmospheric concentrations of silicon dioxide (147) evidenced an increased morbidity rate and a definite lag in general physical development. Rabbits given 2% suspensions of quartz, (176) of particle size 1 to 3 μ and 6 to 12 μ , and aluminum oxide 1 to 3 μ , after 2 to 3 years showed accumulation of the silica particles in different locations according to their size. The largest were found in the pulmonary capillaries, intermediate size in the spleen and hepatic lymph node and the finest ones in the liver. The fine particles were the most active producing nodular cirrhosis of the liver. Coarse particles were less irritating, exciting a simple foreign body reaction which progressed very little in three years. Aluminum oxide particles were merely phagocytosed and produced no fibrosis. A human long term study (006) utilizing slides of the aorta of Japanese patients (known to have a high silica content from talc milled rice in their diet) showed silica in the aorta walls, in the adventitia layer increasing with age, in the intima layer decreasing with age and in the media

Long Term

Studies

remaining constant. At age 40 to 42, the contents of the adventitia and intima layers reversed, perhaps related to hypertension in arteriosclerosis.

Biochemistry

In rats (084,273,511) silica in various forms and in varying amounts was accumulated in body tissue, primarily the liver and spleen in the case of oral administration; the lungs, liver and spleen, when given intratracheally; and the lungs, liver, spleen and kidneys, when given i.v. In the case of colloidal silica administered i.p. there was accumulation of polymerized silica in particle form found in the kidney tubules (041). In the case of humans taking magnesium trisilicate as antacid for extended periods renal calculi with high silica content have been reported (302). A numerical estimation of excretion for rats fed chronic doses (511) showed fecal excretion in the range of 95% of the administered dose, 4% as urinary excretion and 1% accumulated in the tissue. When the silica enters the body through the lungs a considerable amount is caught in the capillary bed of the lungs, some remaining there and some progressively moving to storage sites in other organs (084). When given i.v. the lungs act as a filter for silica particles (359). Renal lesions have definitely been attributed to silica compounds of colloidal particle size (398, 399). Blood protein fractions, histamine (186), and

blood lipids (481) have been notably affected by both crystalline and amorphous silica. Finally, the silica taken up by the macrophages reacts with the membranes surrounding the secondary lysosomes, releasing lytic enzymes into the cytoplasm (012).

Carcinogenicity

Both diatomaceous earth (475) and talc (342) have been shown to produce stomach lesions -- the former in Sherman rats, a rare finding in animals; and the latter in humans on high talc milled rice diet. Also, there are numerous cases of internal granulomas, at times fatal from talc used in surgical procedures (139,213). Locally, widespread granulomata were found as a result of continued heavy use of talcum powder after bathing (496).

Teratology

Colloidal silica (Ludox^R) has been shown to produce retarded lid development or exophthalmia, encephalocele or exencephaly, external edematous blebs, ectopic viscera, retarded feathering and crossed or short upper beak in chick embryos injected in the amniotic cavity (526).

Epidemiology

A high goiter incidence in the high Valle of Aosta (Spain) was attributed to high silica content of the natural water due to runoff from the mountainous and volcanic slopes (493). Also, endemic nephropathy caused by rock erosion in village communities on the banks of the lower reaches of large rivers has been

Consumer

Exposure

reported (329).

The approximate high possible daily intake for the 2-65 year old age group has been estimated by the NAS/NRC GRAS survey to be:

calcium silicate: 481 mg

silica aerogel: 340 mg

sodium aluminum silicate: 2278 mg

sodium calcium aluminosilicate: 2.66 mg

The WHO/FAO monograph on "Toxicological Evaluation of Some Food Colors, Emulsifiers, Stabilizers, and Anti-Caking Agents and Certain Other Substances" (165) set no limit on Acceptable Daily Intakes for the various forms of silicon dioxide, aluminum silicate, calcium silicate, magnesium silicate (including talc) and sodium alumino silicate.

CHEMICAL INFORMATION

I. Nomenclature

Silica (SiO_2) from a number of different sources, and in a number of different forms, is a component of human intake. Forms likely to be assimilated as food additives, and some pure forms occurring in nature which are possible candidates for human intake, are described chemically in Table 1. Throughout this monograph the biological studies described will be considered from the standpoint of the silica, with the form initially administered being specified, when known.

In addition to the forms listed in Table 1, other classifications of silica compounds are considered in the monograph. Oligosilicic acid is a term used to describe mono and lower silicic acids; polysilicic acid refers to the higher, protein-precipitating forms (194). Condensed silicic acids such as disilicic acid ($\text{H}_6\text{Si}_2\text{O}_7$) and trisilicic acid ($\text{H}_4\text{Si}_3\text{O}_8$) occur in rocks which through reaction with the metals therein and by weathering may find their way into the water-supply (262). Diatomaceous earth consists of the siliceous skeletons of the microscopic plant life of the sea - the Diatom. They form the most important food of aquatic animal life (262). This skeleton, or shell, is made up of silica chemically combined with organic and mineral substances (Al, Fe, Ca, Mg, K, Na, H_3PO_4), or the silica may be adhered to a membrane of the Diatom (244). Kahane and Antoine (244), believe that silica participates in a metabolizing system, a state of circulation, before becoming a constituent of the shell of the Diatom, and call this "constitu-

tional silica". The contrasting form, by these authors' definitions, is "interposed silica" which is external in origin and plays its role mechanically as a foreign body.

II. Empirical Formula

See Table 1.

III. Structural Formula

Not Included.

IV. Molecular Formula

See Table 1.

Table 1

Silica Compounds that may be Components of Human Intake *

Common Name	Chemical Name	Formula	Molecular Wt	Description	Melting Point	Solubility
Muscovite	Aluminum Potassium Silicate	$3\text{Al}_2\text{O}_3 \cdot \text{K}_2\text{O} \cdot 6\text{SiO}_2 \cdot 2\text{H}_2\text{O}$	796.40	Monoclinic	Decomposes	Insoluble in cold water
Orthoclase	Aluminum Potassium Silicate	$\text{Al}_2\text{O}_3 \cdot \text{K}_2\text{O} \cdot 6\text{SiO}_2$	556.49	Colorless, monoclinic	1450°C	Insoluble in cold water
Microcline	Aluminum Potassium Silicate	$\text{Al}_2\text{O}_3 \cdot \text{K}_2\text{O} \cdot 6\text{SiO}_2$	556.49	Colorless, triclinic	1150°C	Insoluble in cold water
Sillimanite	Aluminum Silicate	Al_2SiO_5	162.00	White, rhombic	1860°C	Insoluble in cold and hot water and in alkali
	Aluminum Silicofluoride	$\text{Al}_2(\text{SiF}_6)_3$	480.12	White powder		Soluble in hot water, slightly soluble in cold
Zeolex (Trade name)	Aluminum Sodium Silicate	$\text{Al}_2\text{O}_3 \cdot \text{Na}_2\text{O} \cdot 6\text{SiO}_2$	524.29	Colorless, triclinic	1100°C	Insoluble in cold and hot water, soluble in dilute acid
Diopside	Calcium Magnesium Silicate	$\text{CaO} \cdot \text{MgO} \cdot 2\text{SiO}_2$	216.52	White, monoclinic	1391°C	Insoluble in hot and cold water
Pseudowollastonite	Calcium Silicate (α)	CaSiO_3	116.14	Colorless, pseudo-hexagonal	1540°C	Very slightly soluble in cold water, soluble in HCl

Common Name	Chemical Name	Formula	Molecular Wt	Description	Melting Point	Solubility
Wollastonite	Calcium Silicate (β)	CaSiO_3	116.14	Colorless, monoclinic	Transition to α form at 1190°C	
	Calcium Silicide	CaSi_2	96.20			Insoluble in cold water, decomposes in hot water
	Calcium Sili-cofluoride	CaSiF_6	182.14	Colorless		Slightly soluble in cold water, decomposing; soluble in HCl , HF and alkali.
	Calcium Sili-cofluoride (hydrate)	$\text{CaSiF}_6 \cdot 2\text{H}_2\text{O}$	218.17	White, tetragonal		Decomposes in cold water, soluble in H_2SiF_6
	Iron Silicide	FeSi	83.91	Yellow-gray, octagonal		Insoluble in hot and cold water, and in aquea regia
Talc	Magnesium Silicate	$\text{Mg}_3\text{Si}_4\text{O}_{11} \cdot \text{H}_2\text{O}$	379.22	White, monoclinic or rhombic		Insoluble in hot and cold water, and in alkali.
	Magnesium Silicide	Mg_2Si	76.70	Slate blue, cubic	1102°C	Decomposes in cold water and in HCl
	Magnesium Silicide	Mg_5Si_3	205.78		1102°C	Insoluble in cold water; decomposes in hot water, alkali and NH_4Cl

Common Name	Chemical Name	Formula	Molecular Wt	Description	Melting Point	Solubility
	Magnesium Silicofluoride	$\text{MgSiF}_6 \cdot 6\text{H}_2\text{O}$	274.48	Colorless, trigonal	Decomposes	Soluble in hot and cold water.
	Potassium Silicate	K_2SiO_3	154.25	Hygroscopic	976°C	Soluble in cold and hot water, insoluble in alkali
	Potassium Tetrasilicate	$\text{K}_2\text{Si}_4\text{O}_9 \cdot \text{H}_2\text{O}$	352.45	Rhombic	Decomposes at 400°C	Soluble in cold and hot water, insoluble in alkali
Hieratite	Potassium Silicofluoride	K_2SiF_6	220.25	Colorless, hexagonal or cubic	Decomposes	Very slightly soluble in cold and hot water; soluble in HCl; insoluble in alkali and NH_3
Silica Gel	Metasilicic acid	H_2SiO_3	78.08	Amorphous		Insoluble in hot and cold water; soluble in alkali; insoluble in NH_4Cl
Silica Sol	Orthosilicic acid	H_4SiO_4	96.09	Amorphous		Slightly soluble in cold and hot water; soluble in alkali; insoluble in NH_4Cl
	Silicon, crystalline	Si	28.06	Gray, cubic	1420°C, B.P. 2600°C	Insoluble in hot and cold water; soluble in HNO_3 , HF, Ag; slightly soluble in Pb, Zn; insoluble in HF.

Common Name	Chemical Name	Formula	Molecular Wt	Description	Melting Point	Solubility
	Silicon, graphitic	Si	28.06	Crystalline	B.P.2600°C	Insoluble in cold and hot water, soluble in HNO ₃ , HF, fused alkali; insoluble in HF
	Silicon, amorphous	Si	28.06	Brown, amorphous	B.P.2600°C	Insoluble in hot and cold water; soluble in HF, KOH
Opal	Silicon Dioxide	SiO ₂ ·xH ₂ O		Iridescent, amorphous	1600-1750°C sublimates	Insoluble in hot and cold water; soluble in HF, hot alkali, fused CaCl ₂
Cristobalite	Silicon Dioxide	SiO ₂	60.06	Colorless, cubic or tetragonal	1710°C, B.P.2230°C	Insoluble in hot and cold water; soluble in HF; insoluble in alkali
Lechatellierite	Silicon Dioxide	SiO ₂	60.06		B.P.2230°C	Insoluble in hot and cold water; soluble in HF; insoluble in alkali
Quartz	Silicon Dioxide	SiO ₂	60.06	Hexagonal	Transition <1425 B.P.2230°C	Insoluble in hot and cold water; soluble in HF; insoluble in alkali
Tridymite	Silicon Dioxide	SiO ₂	60.06	Trigonal, rhombic	Transition 1670°C	Insoluble in cold water; soluble in

Common Name	Chemical Name	Formula	Molecular Wt	Description	Melting Point	Solubility
					B.P.2230°C	HF: insoluble in alkali
Silica Gel, (Trade name)	Silicon Dioxide, hydrate	$3\text{SiO}_2 \cdot \text{H}_2\text{O}$	198.20	White, amorphous	$-\text{H}_2\text{O}$ at 1200°C	Very slightly soluble in hot and cold water
	Sodium Metasilicate	Na_2SiO_3	122.05	Colorless, rhombic	1088°C	Soluble in cold water, soluble and decomposes in hot water, insoluble in Na or K salts and alkali
	Sodium Metasilicate (hydrate)	$\text{NaSiO}_3 \cdot 9\text{H}_2\text{O}$	284.20	Rhombic	47°C, $-6\text{H}_2\text{O}$ at 100°C	Very soluble in hot and cold water; soluble in .5 N NaOH
	Sodium Orthosilicate	Na_4SiO_4	184.05	Colorless, hexagonal	1018°C	Soluble in hot and cold water
	Sodium Silicate	$\text{Na}_2\text{Si}_4\text{O}_9$	302.23	Amorphous		Soluble in hot and cold water, insoluble in Na or K salts and alkali
	Zinc Silicate	ZnSiO_3	141.44	Hexagonal or rhombic	1437°C	Insoluble in cold water
	Zinc Silico-fluoride	$\text{ZnSiF}_6 \cdot 6\text{H}_2\text{O}$	315.54	Colorless, hexagonal	Decomposes	Soluble in cold and hot water.

* Data extracted from Lange's Handbook of Chemistry.

V. Specifications

A. Aluminum Sodium Silicate (Zeolex)

1. FDA specifications, data supplied by them.

Procedure: Codex

Silicon Dioxide: Not less than 66%, not more than 71% of SiO_2 after drying.

Al_2O_3 : Not less than 9%, not more than 13% of Al_2O_3 after drying.

Na_2O : Not less than 5%, not more than 6% of Na_2O after drying.

Loss on drying: Not more than 8%

Heavy Metals:

Arsenic as As Not more than 3 ppm (0.0003%)
Heavy Metals (as Pb) Not more than 10 ppm (0.001%)

Must meet Codex specification.

2. Food Use - Commercial specifications (data supplied by the FDA).

Color and Appearance White Powder.

Granulation
On USBS Sieve #100 0.2% Maximum.
On USBS Sieve #200 0.5% Maximum.
On USBS Sieve #325 1.0% Maximum.

Moisture 9.0% Maximum

Odor Clean, free from objectionable odors.

pH 10.0 ± 1.0

Visual Impurities None

B. Calcium Silicate

1. Commercial specifications - Technical Grade for industrial use only (paint, rubber and plastics industries). Data supplied by the FDA.

CALCIUM SILICATE (1) $3\text{CaO} \cdot 3\text{SiO}_2$

<u>Item</u>	<u>Typical Analysis</u>	<u>Specification</u>
SiO ₂	66%	
CaO	19%	
NaCl	1.43%	2.0% max.
Loss at 105°C.	2.1%	6.0% max.
Ignition Loss above 105°C.	9.7%	
R ₂ O ₃	0.7%	
MgO	0.2%	
Cu and Mn (Total)	0.004%	
As	1 ppm	
Hg	< 0.1 ppm	
Pb	6 ppm	
F	40 ppm	
Heavy Metals as Lead	15 ppm	

2.

CALCIUM SILICATE (2) $\text{CaO} \cdot 12\text{SiO}_2$

<u>Item</u>	<u>Typical Analysis</u>	<u>Specification</u>
SiO_2	80%	83% max.
CaO	4.0%	7% max.
NaCl	1.0%	2% max.
Loss at 105°C.	6.0%	3-9%
Ignition Loss above 105°C.	8.0%	8% max.
Fe_2O_3	0.13%	0.20% max.
Al_2O_3	0.45%	1% max.
R_2O_3	0.60%	
As	0.3 ppm	1 ppm max.
Hg	0.1 ppm	1 ppm max.
Pb	8 ppm	10 ppm max.
F	7 ppm	50 ppm max.
Heavy Metals as Lead	25 ppm	25 ppm max.

C. Silicon Dioxide

1. Commercial specifications, for industrial use in the rubber and plastics industries. Data supplied by the FDA.

SILICA - HYDRATED, PRECIPITATED

<u>Item</u>	<u>Typical Analysis</u>	<u>Specifications</u>
SiO ₂ (dry basis)	94%	87.0% minimum
pH in 5% Water Suspension	7.0%	6.5 - 7.3
Loss at 105°C.	5.3%	3.0 - 7.0%
NaCl	1.7%	
CaO	0.80%	
R ₂ O ₃	0.63%	
Combined Cu and Mn	0.003%	
As	0.5 ppm	
Hg	<0.5 ppm	
Pb	5 ppm	
Heavy Metals as Pb	15 ppm	

2. Food Use - as a conditioning agent - commercial specification.

Data supplied by the FDA.

SILICON DIOXIDE

Color and Appearance	White powder.
Moisture	1.5% Maximum.
Odor	Clean, free from objectionable odors.
pH	3.9 ± 0.4
Visual Impurities	None.
Loss on Ignition	3.0% Maximum.
Bulk Density, Tapped	0.071 ± 0.006

3. Food Chemicals Codex, 1972 Edition (104).

Description

Silica aerogel is a white, fluffy, powdered or granular micro-cellular silica. Hydrated silica is a precipitated, hydrated silicon dioxide occurring as a fine, white, amorphous powder, or as beads or granules. Both forms of silicon dioxide are insoluble in water and in alcohol and other organic solvents, but are soluble in hot phosphoric or hydrofluoric acids and in solutions of alkalis at 80° to 100°.

Specifications

Assay

Silica aerogel: Not less than 90.0% of SiO_2 .

Hydrated silica: Not less than 89.0% of SiO_2 , calculated on the dried basis.

Loss on drying. Hydrated silica: not more than 6%.

Loss on ignition. Not more than 6%.

Limits of Impurities

Arsenic (as As). Not more than 3 parts per million (0.0003%).

Heavy metals (as Pb). Not more than 3 parts per million (0.003%).

Lead. Not more than 10 parts per million (0.001%).

Soluble ionizable salts (as Na_2SO_4). Not more than 5%.

Tests

Assay. Transfer about 2 grams, accurately weighted, into a tared platinum crucible, ignite at 600° for 2 hours, cool in a desiccator, and weigh. Moisten the residue with 7 or 8 drops of alcohol, add 3 drops of sulfuric acid, and add enough hydrofluoric

acid to cover the wetted sample. Evaporate to dryness on a hot plate, using medium heat (95-105°), then add 5 ml. of hydrofluoric acid, swirl the dish carefully to wash down the sides, and again evaporate to dryness. Ignite the dried residue to a red heat over a Meker burner, cool in a desiccator, and weigh. The difference between the total weight loss and the weight loss after ignition at 600° represents the weight of SiO_2 in the sample taken.

Loss on drying, page 931. Dry hydrated silica at 105° for 2 hours.

Loss on ignition. Transfer about 5 grams, accurately weighed, into a suitable tared crucible, and ignite at 600° to constant weight.

Sample Solution for the Determination of Arsenic, Heavy Metals, and Lead. Transfer 10.0 grams of the sample into a 250-ml. beaker, add 50 ml. of 0.5 N hydrochloric acid, cover with a watch glass, and heat slowly to boiling. Boil gently for 15 minutes, cool, and let the undissolved material settle. Decant the supernatant liquid through a Whatman No. 3 filter paper, or equivalent, into a 100-ml. volumetric flask, retaining as much as possible of the insoluble material in the beaker. Wash the slurry and beaker with three 10-ml. portions of hot water, decanting each washing through the filter into the flask. Finally, wash the filter paper with 15 ml. of hot water, cool the filtrate to room temperature, dilute to volume with water, and mix.

Arsenic. A 10-ml. portion of the Sample Solution meets the requirements of the Arsenic Test, page 865.

Heavy metals. Dilute 20.0 ml. of the Sample Solution to 30.0 ml with water. A 10-ml portion of the dilution meets the requirements of the Heavy Metals Tests, page 920, using 20 mcg. of lead ion (Pb) in

the control (Solution A).

Lead. A 10-ml. portion of the Sample Solution meets the requirements of the Lead Limit Test, page 929, using 10 mcg. of lead ion (Pb) in the control.

Soluble ionizable salts. Weigh accurately 12.5 grams of the sample, and stir it with 240 ml. of water for at least 5 minutes with a high speed mixer. Transfer the mixture into a 250-ml. graduate, and wash the mixer container with water, adding the washings to the graduate to make 250 ml. Stopper the graduate, and invert it several times to mix the slurry. The conductivity of the slurry, determined with a suitable conductance bridge assembly; is not greater than that produced by a control solution containing 750 mg. of anhydrous sodium sulfate in each 250 ml.

Packaging and storage. Store in well-closed containers.

Functional use in foods. Anticaking agent; defoaming agent.

VI. Description

See Table 1.

VII. Analytical Methods

1. For the determination of silica in acid-insoluble silicates such as talc or kaolin, McClellan (336) offers a gravimetric method which yields a recovery of ca 99.6%. The procedure utilizes a strong acid for dissolving the fusion melt and dispenses with evaporation for dehydrating the silica. A sample is fused with sodium carbonate (in a platinum crucible at 950-1050°C for 15 minutes. The melt is then dissolved with HNO_3 which precipitates about 95% of the silica of

the sample. This suspension is treated with NH_4Cl , HClO_4 and H_2SO_4 , and heated for 15 minutes after the evolution of oxides of nitrogen has ceased. The cooled mixture is then diluted with hot water, filtered, and ignited, with the precipitate being weighed as crude silica. Correction for impurities is made by volatilizing the silicon as silicon tetrafluoride.

2. King, Stacy, Holt, Yates and Pickles (257) point out that all analytical methods for silica are based on the decomposition of siliceous material to liberate silicic acid. They have thoroughly reviewed prior work in some detail (257), and have undertaken a critical investigation of the basic ferric acetate precipitation procedure used for removing phosphate from solutions prior to silica determination. They find that unless there is strict control of pH, inaccurate results may be obtained, due to incomplete phosphate removal or loss of silica with the phosphate precipitation. Also they confirmed previous work that showed both phosphate and silica couple with molybdate in weak acid, but that only the silicomolybdate complex is reduced in strong acid. This principle is the basis for their procedure for determining silica in biological material, with the phosphate left in solution. Iron, in amounts greater than those occurring in animal tissue, does not interfere. In urines with high phosphate concentrations they have devised two procedures to overcome interference by precipitation of ammonium phosphomolybdate. Phosphate may be partly or completely removed by precipitation with nitrous acid and sodium molybdate is used to couple with silica.

Their colorimetric procedure agrees well with gravimetric analy-

ses, and good recoveries (in these authors' estimation) of added silica have been obtained from tissue, blood, and urine. The method was also tested on small samples of mineral dusts; results agreed well with those obtained with gravimetric methods.

3. Further attention to the phosphate interference in analysis for silica in urine has been given by Paul (384). His procedure involves precipitating the bulk of the phosphate by calcium chloride and ammonia at pH 7, and then removing the remaining phosphate by selective extraction of the phosphomolybdate with ethyl acetate. Silica is then determined in the aqueous phase after reduction of the silicomolybdate by 1-amino-2-naphthol-4-sulphonic acid in the presence of citric acid. The author indicates the method is accurate and reproducible, and generally applicable to trace amounts of silica in high phosphate concentrations.

4. Fast-neutron activation analysis has been utilized for the determination of silicon in sputum (430). The procedure was developed to identify and separate cases of silicosis from those of pulmonary tuberculosis. Sputum is digested by trypsin, centrifuged, and heated with trichloroacetic acid to remove phosphorus. For irradiation, a neutron generator with a fast-neutron yield of 2×10^9 n per cm^2 per second is used. After an exposure of 2-3 minutes, followed by a one-minute delay, the activity is measured with a multi-channel analyzer, for 4-6 minutes. The silicon content is evaluated from the aluminium-28 activity produced by the reaction $^{28}\text{Si}(n,p) ^{28}\text{Al}$ with a photopeak at 1.78 MeV. Although fast-neutron activation analysis by use of neutron generators had not been applied

to biological substances at the time of this author's publication, he points out that the method seems promising for such analyses.

5. Continuous on-stream instrumentation is finding increased application in control analysis, and Sheen and Serfass (451) have described its application to silica determinations. In this analysis, four specific reagents are metered to the zero and sample cells of the analyzer. First a buffer is metered to maintain pH as close as possible to 1.7. Ammonium molybdate, which forms heteropolymolybdic acids with the silica and phosphorus in solution, is also metered. The complexing agent then is metered to both cells and attacks the phosphomolybdic acid, destroying it completely. The reagent has no effect on the silicomolybdic acid. Finally a reducing agent is added, resulting in the formation of colloidal molybdenum blue. This reaction always goes to completion, and therefore the color intensity is a direct measure of the soluble silica present. For the blank cell comparison, chemical reagents are fed to a cell to complete their reactions before being fed to the zero cell. Each complete cycle normally lasts 12 minutes. The range of the instrument is normally 0 to 50 ppb, and a change of 1 ppb in water may be detected accurately.

VIII. Occurrence

Hydrophile polymeric silicic acids are used as food additives, to stabilize emulsions, as antisedimentation agents for cocoa in beverage mixes, as dispersing agents, and as clarifying agents for beer (291). Talc (magnesium silicate) is used extensively in Japan as a milling agent for rice (342). According to the NAS/NRC Compre-

hensive Gras Survey Report (375) calcium silicate, silica aerogel, and aluminum sodium silicate each are used in as many as 14 different categories of food, as stabilizers and emulsifiers. Talc (hydrous magnesium silicate) is used as a filler in oral pharmaceutical preparations (542) and magnesium trisilicate as an antacid (302 and others).

In plants, the role of silicon seems to correspond to that of calcium and phosphorus in animals (262). One family of sponges has a skeleton of almost pure silica; the ash of wheat straw may contain 40% SiO_2 ; and the equisetum (horse tail) has been shown to contain 80% SiO_2 (262). Another indication of the high silica content of plants is its increased occurrence in tissues of herbivorous animals as compared with that of carnivorous animal tissues (262).

Silicon seems invariably to be present in animal matter; also all marine forms contain silica (262). All tissues and fluids examined contain at least traces, and in many cases amounts are within the range of those of the mineral elements recognized as being normal tissue constituents (262). King and Belt (262) conclude that the almost universal distribution of silica makes it probable that silicon is a likely contaminant, if not an essential ingredient, of all protoplasm. Their data on silica in a variety of types of tissue are given in Table 2.

Table 2
Silica Content of Normal Tissues (262)

FETAL TISSUES	SiO ₂ DRY TISSUE <i>mgm. per 100 grams</i>	ADULT TISSUES (HUMAN)	SiO ₂ DRY TISSUE <i>mgm. per 100 grams</i>
Whole mouse ⁷	10	Heart: myocardium ¹	11-29
Blood (human) ⁸	13	Heart: endocardium ¹	14-28
Brain (human) ⁷	22	Ileum ¹	47
Heart (human) ⁷	20	Kidney ¹	11-27
Kidney (human) ⁷	13	Liver ^{7, 1}	11-17
Kidney (calf) ⁷	8-10	Lung ^{7, 8, 9, 1} (infancy to old age)	10-200
Liver (human) ⁷	4-6	Lung ¹ (infancy to old age)	14-2000
Liver (calf) ⁷	4-10	Lymph glands ¹ (peribron- chial)	27-5000
Lung (human) ⁷	8-10	Mammary gland ¹	12
Lung (calf) ⁷	4-10	Milk ^{8, 10}	71
Lung (human) ⁸	31	Muscle: pectoral ¹	15-15
Muscle (calf) ⁷	21-37	Muscle: psoas ¹	23
Spleen (calf) ⁷	11	Nerve: peripheral ¹	11-40
ADULT TISSUES (HUMAN)		Nerve: spinal cord ¹	13-87
Adrenal ¹	14-80	Ovary ¹	10-21
Appendix ¹	68	Oesophagus ¹	8-26
Aorta thoracic ¹	11-31	Pancreas ^{1, 3}	14-31
Artery iliac ¹	14-50	Parathyroid ¹	22-23
Bladder ¹	7-21	Pituitary ¹	27-67
Bone ^{7, 11, 3}	12-30	Placenta ⁸	29
Brain: dura mater ¹	11-37	Prostate ¹	11
Brain: cortex ¹	12-36	Rectum ¹	15-74
Brain: Cerebellum ¹	16	Skin: scalp ¹	35-270
Brain: pons ¹	13-43	Skin: abdomen ¹	40-130
Brain: pineal ¹	70-90	Skin: sole of foot ¹	11-59
Cecum ¹	20-27	Spleen ¹	15-41
Cartilage: costal ¹	18-35	Stomach ¹	12-12
Cervix ¹	11	Teeth ²	6
Diaphragm ¹	11-50	Tendon, Achilles ¹	12-16
Duodenum ¹	10-80	Thymus ¹	81
Gall bladder ¹	31	Thyroid ¹	21-55
		Testes ¹	8-20
		Umbilical cord ¹	33
		Uterus ¹	7-12

The most important constituents of geological formations are the silicates. As Belt and King have pointed out (262), whole mountain ranges are composed of them. The natural silicates found in rocks, correspond to the disilicic and trisilicic acids (condensed silicic acids mentioned in Section I). For example, wollastonite (CaSiO_3) is a meta silicate, willemite (Zn_2SiO_4) an ortho silicate, serpentine ($\text{Mg}_3\text{Si}_2\text{O}_7$) a disilicate, and orthoclase (KAlSi_3O_8) a trisilicate. In the weathering process carbon dioxide dissolved in natural waters attacks those silicates which contain metals capable of combining with carbonic acid, and colloidal silica is liberated (262). with the metal silicate being left as residue. \(\backslash\) Trikurakis (493) reported on a study of the goiter incidence in the high Valle of Aosta (Spain). This was attributed to high silica content of the natural water due to runoff from the mountainous and volcanic slopes, with the condition being strongest in the lowest parts of the mountain slopes.

As noted previously, silica is found in all natural waters -- mineral water, hot springs, city water, well-water and sea water. Akiya, et al., (006) collected samples of water in Japan, Europe, North and South America, and Southeastern Asia. These samples were collected in polyethylene bottles and returned to the home laboratory for analysis. Results are shown in Table 3.

Table 3

SiO₂ and CaCO₃ Contents in Water of America, Europe and Southeastern Asia. (006)

Place	Country	Sorts of water	Free Si as SiO ₂ (ppm)	Total Si as SiO ₂ (ppm)	Hardness CaCO ₃ (ppm)	Remark
San Francisco	U. S. A.	Drinking water		2.28	32.07	City water
São Paulo	Brazil	"		2.89	45.82	"
Campinas	"	"		18.15	51.97	"
Frankfurt	Germany	"		7.33	32.07	"
Hamburg	"	"		3.91	227.91	"
Paris	France	"		4.92	265.71	"
Basel	Switzerland	"		4.64	100.79	"
Teheran	Persia	"	8.0	19.2	176.0	Well water
Karachi	Pakistan	"	6.0	8.0	100.0	City water
Bombay	India	"	7.8	26.1	25.2	Lodging House
"	"	"	17.8	42.4	590.0	Khora-village
"	"	"	19.4	29.2	250.0	"
Culebra	"	Bath water	16.2	34.4	460.0	Grand Hotel
"	"	Drinking water	17.2	22.0	430.0	"
"	"	"	6.8	13.8	72.8	Toyo Menka Co.
New Delhi	"	"	3.9	80.0	172.0	City water
Rangoon	Burma	Lake water	5.9	13.3	16.3	Hlawga Lake
Singapore	Singapore	Drinking water	6.4	9.1	28.5	City water
Jakarta	Indonesia	"	8.6	21.3	32.9	"
Bangkok	Thailand	"	6.4	17.6	36.2	"
Phnom Penh	Cambodia	"	13.1	16.6	63.0	"
Saigon	Vietnam	"	8.6	16.8	12.0	"
Manila	Philippine	"	24.1	29.2	36.0	"

Akiya (006) also reported that Pacific Ocean sea water contains only from 0.5 to 3 ppm of silica; in general the city water of Europe and America contains from 2 to 7 ppm.

Atmospheric dust in the vicinity of factories manufacturing silicon and its alloys has been reported to contain 30 to 66% of free silica; mean concentrations of such dust of up to 3.1 mg/m^3 have been reported by Egorova (147).

IX. Stability in Containers

A study of the influence of storage time on the adsorption properties of silica zerogels has been made by Sheinfain and Stas (452). Four different zerogels were prepared from hydrogels, each with different adsorption and structural characteristics, as determined by measuring the isotherms of methyl alcohol vapor adsorption at 20°C . The samples were stored for four years in loosely stoppered jars, and methyl alcohol vapor isotherms again measured. It was found that, regardless of original structure, the prolonged storage led to decreased adsorption over the entire range of relative pressures of the adsorbate. Specifically, the data showed that the pore volume and specific surface decreased during aging, giving an increase in structure density. The authors suggested that the changes were due to the action of atmospheric moisture.

BIOLOGICAL DATA

I. Acute Toxicity

A. Animal Studies: (see Table 4).

Table 4

Substance	Animal	Sex and Number	Route	Dosage mg/kg Body Wt	Measurement	Reference
Freshly prepared Silica Sol	Mice		i.p.	50-100	LD	209
Silica Sol, opalescent and somewhat gelled	Mice		i.p.	250	LD	209
Fresh Silica Sol (ca 3%)	Mice		i.v.	100	LD (in a few minutes)	209
Fresh Silica Sol (ca 1%)	Mice		i.v.	150	LD (in 3 minutes)	209
Silica Sol, older opalescent 1%	Mice		i.v.	50	LD (in 3 minutes)	209
Silica Sol, fresh	Mice		i.v.	100	LD (2-24 hours)	209
Soluble Silica in Saline	Mice	6	i.v.	100	LD (24 hours)	209
Fresh (1/2 hr old) SiO ₂ (ca 2mM)	Mice		i.v.	180 (fractionated)	LD	194
Solution above after 80 days	Mice		i.v.	9	LD	194

Table 4 (Cont'd)

Substance	Animal	Sex and Number	Route	Dosage mg/kg Body Wt	Measurement	Reference
Synthetic amorphous Silica (av particle size 0.01 μ)	Mice		i.v.	9	LD	194
Talc	Mice		i.p.	4250	LD	056
Fresh Silica Sol (ca 3%)	Rats		i.v.	100	LD (in a few minutes)	209
Precipitated Silica, particle size 19 μ	Rats		i.v.	35.2	LD ₅₀ (in a few minutes)	084
Fresh precipitated Silica, particle size 20 μ	Rats		i.v.	41.2	LD ₅₀ (in ca 2 hours)	084
Precipitated Silica, particle 25 μ	Rats		i.v.	44.4	LD ₅₀ (in ca 2 hours)	084
Silica Condensate 150 μ particle size suspended in isotonic saline	Rats		i. trach.	30 & 50	LD ₈₀₋₉₀ (in a few hours)	397
Same solution as above	Rats		i.p.	100	LD ₂₀₋₃₀	397
Fresh Silica Sol	Guinea Pigs		i.p.	120	LD	209
Silica Sol, opalescent and somewhat gelled	Guinea Pigs		i.p.	200-240	LD	209
Fresh Silica Sol (ca 3%)	Guinea Pigs		i.v.	100	LD (in a few minutes)	209

Table 4 (Cont'd)

Substance	Animal	Sex and Number	Route	Dosage mg/kg Body Wt	Measurement	Reference
Fresh Silica Sol* (ca 3%)	Rabbits		i.v.	100	LD (in a few minutes)	209
Silica, 1% in isotonic saline	Rabbits		i.v.	141	LD (3 minutes, blood clot in right side of heart)	209
Silica Sol, opalescent, 3%	Rabbits		i.v.	35	LD (2 minutes, blood clot in right side of heart)	209
Soluble silica, 1.4%	Rabbits		i.v.	75 (in 7 doses, ca 1/2 hr apart)	LD (immediately after last injection)	209
Silicic Acid, 8 batches similarly prepared, but with pH varying from 1.53 to 2.65	Cats	53	i.v. (Rate of inj. varied)	15-193	Av LD	348
Colloidal Silicic Acid	Dogs	3	i.g.	25-50 cc	LD (3 days)	476
Particulate Silica (Quartz powder in isotonic NaCl)	Dogs	1	i.v.	20	LD (6 hours)	260

* The authors concluded from the nature of variation that there was no correlation between LD and pH, or between LD and injection rate. However, within each batch the LD remained in the same general range.

B. Humans

1. A fatal case of talc aspiration has been reported by Molnar, Nathenson and Edberg (351). A 22-month-old Negro boy was found by his mother, choking, breathing heavily, and coughing, with a talcum powder can beside him, and talc in his mouth. Vinegar and milk failed to induce emesis, and the child was rushed to the hospital. Initial examination indicated moderately severe respiratory distress with perioral cyanosis. The mouth and pharynx were clear; there were bilateral coarse rhonchi in the chest, with grunting; but the heart sounds were normal. Although he was given penicillin and chloramphenicol and placed in an oxygen tent with cool mist after 7 hours his respiratory distress increased. By this time bilateral moist rales and rhonchi had developed throughout both lung fields. He was then treated with lanacide C, morphine, and meralluride, for cardiotonic effect. There was some improvement, but intractable cardiopulmonary failure gradually developed and the child died 20 hours after talc aspiration. At autopsy the oral cavity, pharynx, trachea and esophagus were free of foreign material. However, the mucous membranes of the larynx, trachea and main bronchi were covered with viscid, transparent mucus and were slightly injected. The lungs were voluminous with pink and dark red segments. The pleural surfaces were free of any exudate or nodules. A greenish-yellow creamy material filled small bronchi on the lung surface, and all organs exhibited marked congestion. Microscopically, all sections of the lungs showed focal areas of edema and emphysema, and under polarized light the distribution of talc could clearly be seen.

They were present in minute quantities, in sizes from 2 to 50 μ in length and 1 to 10 μ in thickness. Intracytoplasmic particles of talc could be detected in the leukocytes and macrophages. The post-mortem diagnosis was acute bronchitis and bronchiolitis due to aspiration of talcum powder.

A near fatal case, of talc aspiration which followed a similar course, up to the time when dexamethasone was initiated, has been reported by Hughes and Kalmer (233). The authors also cited two other fatal cases, in addition to the death described above.

2. Fatalities due to talc embolism have only recently been recognized (542). Such cases arise following intravenous use of crushed drugs, compounded for oral therapy and containing talc, by "main liner" drug addicts. Zientara and Moore (542) describe the results of an autopsy on a 31-year-old woman who had been a heroin addict since the age of 16. She was admitted to the hospital with acute respiratory failure, and died within minutes. Gross examination indicated heavy, stiff lungs with multiple hard white nodules scattered throughout the lung parenchyma; in some areas they became confluent to form larger masses. The heart was dilated with slight hypertrophy of the right ventricle. Microscopically the lungs showed talc particles in both pulmonary arteries and in surrounding interstitial tissues, causing foreign body granulomas. The walls of the pulmonary arteries were thickened, with angiomatoid lesions, and thrombi related to talc particles were present. Similar particles were also found in the liver, spleen and kidney. The authors did not know the exact source of

the talc emboli but suspected that it was derived from methadone tablets, injected intravenously in suspension.

3. Atlee (043) has reported a characteristic retinopathy due to talc and cornstarch emboli in 17 drug addicts. The occurrence followed repeated i.v. injections of crushed methylphenidate hydrochloride tablets taken for stimulation. Ophthalmoscopic examination showed tiny glistening crystals, mostly in the small blood vessels around the macula. Two of the patients, who had pulmonary hypertension, had reduced vision assumed to be caused by retinal edema. Histological examination of tissues from one patient confirmed the presence of talc and cornstarch in the retina and choroid, as well as in the lungs and brain.

II. Short Term Studies

A. Rats

1. Diatomaceous earth, the siliceous remains of unicellular and colonial algae, is used in insect control in granaries, and the grain so treated will have a residue which could conceivably be assimilated by humans (061). Bertke (061) in an effort to determine possible tissue damage studied the effects on rats over a 90-day period. Weanling white rats, Wistar strain (av wt 50 g), 15 males and 15 females were fed diatomaceous earth mixed as a 5% water suspension in their feed. Weekly weight changes were recorded, and the animals were sacrificed at 90 days. Autopsies were performed and histologic sections were made from the stomach, small and large intestines, liver, kidney, spleen, lung,

urinary bladder, adrenal glands, mesenteric lymph nodes, and testes or ovaries. Parts of the liver, kidney, and spleen were analyzed for residual silica. Two other groups of the same size received 3% and 1% respectively in their diets, with the same experimental and examination protocol followed; and the untreated group IV rats were examined as controls. Particle-size distribution measurements of the test material showed, that 90% of the particles were <0.1 mm in diameter; 55% were <0.012 mm; the smallest was 0.00046 mm and the largest 0.64 mm.

The average weekly weight gain of the 5% group was higher after the first week than the gain of the controls; the maximum weight gain differential was reached during the sixth week. In following weeks, although test animals continued to gain more rapidly than controls, the differential became progressively smaller. Projection beyond the 90 day period indicated that weight gains of the two groups possibly would be equal by the 14th or 15th week. No accurate averages were undertaken on the lower dose groups. Complete histologic studies were made on the 5% group; six rats each from the 3% and 1% categories were studied for comparison. No differences were observed in sections of these test animals when compared with sections from the controls. Silica accumulation in liver, kidneys, and spleen was negligible.

2. In another attempt to determine the damage caused by oral ingestion of silica particles McClendon et al., (337) treated six groups of weanling rats, 10 per group, with silica in the forms and of the particle sizes shown in Table 5.

These rats were fed 90% Purina Dog Chow and 10% of one of the forms listed for 3 months. A seventh group of ten untreated rats was used as a control. After sacrifice, histological examinations were made of the stomach, small and large intestine, liver, spleen, pancreas, adrenals, mesenteric lymph nodes and lung. As in the case of diatomaceous earth treatment no lesions were found in any of the organs. No analysis was made for residual silica in the organs.

3. Kochmann and Maier studied the effects of four different silicic acid preparations (273). Groups of almost fully grown albino rats, five each, were fed diets containing bread and dried milk. The first group additionally received 15 cc of distilled or mineral water containing silicic acid. The other groups were fed the same diet with the addition of commercial silicic acid preparations, i.e. Siliquid (a colloid silicic acid solution with 0.25% SiO_2), colloid silicic acid with 1% SiO_2 , or Silistren (an O-silicic acid tetra glycolester with 18 to 20% SiO_2). Each of the commercial preparations contained the same amount of silicic

Table 5.
Forms and Particle sizes of Silica Fed to Weanling Rats (337)

Name	Particle size in μ
Linde silica	.02-.05
Columbia-Southern Hi-Sil 101	.03
Dow Corning silica (unpelletized)	.01-.02
Powdered silica gel	
Sodium metasilicate	
Carborundum Co. Fibrofrax (silica fibers)	

acid as the same volume of natural mineral water. Tables 6 and 7 below indicate that the total SiO_2 for the mineral water was 50 mg, and for the commercial preparations 70 mg each. Growth values are also shown.

Table 6

Experimental Series of 57 Days' Duration (273)

	Average weight of 5 rats each	
	with 15 cc H_2O	with 15 cc Silvana hot spring water
Initial weight	151 g	141.5 g
Final weight	181 g	182 g
Increase	30 g = 20%	40.5 g = 29%
Daily increase	0.53 g	0.71
Ratio of increase	100	145
SiO_2 total		50 mg

Table 7

Experimental Series of 60 Days' Duration (273)

	Average weight of 5 animals each			
	Control Animals	Siliquid	Colloid SiO_2	Silistren
Initial weight ...	85.0	94.0	90.0	92.0
Final weight	148.4	157.8	152.8	139.3
Increase	63.4 = 74%	63.8 = 68%	62.8 = 70%	47.3 = 51%
Daily increase ...	1.06	1.06	1.05	0.79
Ratio of increase	100	92	95	69
SiO_2 total		70 mg	70 mg	70 mg

The authors point out, on the basis of these data, that the animals receiving mineral water showed an increase in growth rate over that of controls. This was not the case, however, with the groups fed the commercial preparations. Animals fed on Siliquid and colloidal silicic acid had similar growth rates to the controls, while the Silistren group showed a pronounced lag in body weight increase that the authors indicated was beyond the limits of experimental error. No theory was offered for this difference, other than that the silicic acid might be in some special combination that inhibited growth. They did postulate that the increased growth rate of the mineral water group was probably due to the natural components of the water, i.e. iron, carbon dioxide etc. Accumulation of silica will be discussed later.

4. Still another investigation of the ingestion of silica in rats was conducted by Webb, Selle, and Thienes (511). They undertook the study to test the possibility that high silica content of table salt might be the etiological factor in the high silica content of eye lens and resulting cataract, as noted in India. This correlation of lens silica content with cataracts has not been found in American cases. Four series of rats were studied: Series I - controls, Series II 0.01% SiO_2 (as Braun's Air Float, 325 mesh) in their food, Series III 0.1% SiO_2 , and Series IV 1.0% SiO_2 . The animals were on this regime for 10 months, and then sacrificed and the tissues prepared and analyzed for silica. Eyes of none of the rats showed any pathological change in the lens or other media for transmitting or refracting light. No detrimental

changes were observed in the animals during the course of the experiment, and growth rates for all four series were the same. Some silica accumulated in the liver, spleen and testis -- particularly in series III and IV -- but lens silica values in the three test groups corresponded, within experimental error, to that of the controls. Accumulation in the various organs is discussed in more detail in Section II of Biochemical Aspects.

5. Byers and Gage (084) investigated the toxicity of "precipitated silica" -- i.e. amorphous silica prepared by acid precipitation from an aqueous solution of sodium silica in a process similar to that used for the preparation of silica gel. This is a commercially important type, according to these authors, and they undertook their investigation to assess the possibility of a silicosis hazard from its industrial use. This may not seem particularly pertinent to the evaluation of silica compounds as food additives; nevertheless, some information regarding the action of silica from such a source in the animal body may be of value. Samples from two suppliers were studied. From supplier A came A1, of particle size 19 μ as calculated by an air permeability method; A2 of particle size 20 μ when freshly prepared, as shown by electron microscopy, and at a later date 60 μ as indicated by air permeability measurement. Supplier B provided Sample B of particle size 25 μ , again as measured by air permeability. Sample A2 particles seemed to have a greater tendency to aggregate, when tested in dust cloud chamber than did those of either A1 or B. Also there was greater difficulty encountered in preparing an

aqueous suspension of A2. The acute toxicities of these three samples by i.v. injection were reported in Table 4.

Groups of 50 M and 50 F adult Wistar albino rats were injected intratracheally, under light ether anesthesia, with 1 ml of a 2.5% suspension of each sample. The silica had been sterilized in hot air and dispersed in sterile isotonic saline. Higher doses than 25 mg/rat caused death within 24 hours. Twelve rats from group A2 died within 4 days, presumably from lung infection since this group was not injected with penicillin as groups A1 and B had been. From each group 5 M and 5 F were sacrificed at three intervals during the 12 months of the study, and lungs, liver, kidneys, and spleen were taken for silica determinations. Similar groups were also sacrificed for histopathological examination of the lungs. A significant amount of silica was detected in the lungs after 12 months, but a rate of elimination much faster than that for quartz and other fibrogenic dusts was noted. Lesions were produced, mild fibrosis, but again the authors stress they did not resemble those from quartz. Sample A2 produced the most marked effect, with the lesions the largest and most persistent, and with a slightly greater deposition of fibers than with A1 and B. The lesions with all three groups regressed during the course of the experiment, corresponding to the disappearance of silica from the lungs. The authors evaluate the lung function effect of the three dust samples as small. There was evidence of mild emphysema in a few lungs but aside from a few large abscesses and foci of bronchiectasis there was little evidence of infection.

The authors conclude that the dusts did not greatly influence the incidence or course of infections, when compared to lung function of normal controls. Numerical data and detailed discussion of accumulation and excretion are discussed in a later section.

6. Silica and several silica salts were studied by Newberne and Wilson (372). Groups of 15 M and 15 F Charles River CD strain rats (BW 80 to 100 g) were fed silicon dioxide, 0.8 g/kg/day; aluminum silicate, 1.3 g/kg/day; sodium silicate, 2.4 g, or magnesium trisilicate, 1.8 g, in a semisynthetic diet. After four weeks the animals were sacrificed. The only clinical symptoms during the feeding period were polydipsia, polyuria, and soft stools, seen intermittently in a few animals fed magnesium trisilicate or sodium silicate. All clinical chemical tests were within normal limits. No drug-related lesions occurred in any of the rats, and the only observed departure from normality was an occasional rat from each group with an isolated hyaline tubular cast. Dogs treated similarly showed gross microscopic renal lesions from sodium and magnesium silicates, to be discussed in a later section.

B. Guinea Pigs

The mineral talc is a specific substance, anhydrous magnesium silicate $[\text{H}_2\text{Mg}_3(\text{SiO}_3)_4]$. Several minerals may accompany it as it occurs in nature and some of them may be quantitatively more important than talc. Most prominent are serpentine (hydrous magnesium silicate, $[\text{H}_4\text{Mg}_3(\text{SiO}_3)_4]$), dolomite (calcium magnesium carbonate) and tremolite (calcium magnesium silicate). Talcs used in industry (in cosmetics, as fillers in paper, paints, as dusting

agents, etc.) vary greatly. Also it is known that talc dust exposure results in diffuse fibrosis. Schulz and Williams (441) studied the effects on guinea pigs of seven commercial talcs, choosing their samples to represent the range of composition, with regard to the $\text{Si}:\text{CO}_2$, that was covered by 51 different talcs they had analyzed. Table 8 reports the composition of the 7 samples tested.

Table 8
Talc Samples (441)

	WT. % SOLUBLE	TALC	SERP.	CARD.	QTZ.	TRM.	OTHERS	Si:CO ₂ RATIOS
		%	%	%	%	%	%	
1. 814-6A!	41.73	6	5	86	—	—	3	15:85
2. 814-11C	35.56	12	35	45	tr	tr	8	55:45
3. 814-25G	22.94	16	26	47	tr	tr	11	53:47
4. 814-33	14.68	24	30	32	tr	8	6	68:32
5. 814-40B	7.95	32	49	9	tr	tr	10	91:9
6. 814-50	1.13	52	39	—	—	—	9	100:0
7. 814-34	12.52	3	3	8	tr	82	4	92:8

Guinea pigs (BW 250 g), in groups of 10, were injected once intraperitoneally (i.p.) with a 4-cc suspension (5 g in 100 cc physiological saline) having the equivalent of 200 mg of talc. They were sacrificed at intervals from 10 days to 15 months, and tissues from the ventral parietal surface of the peritoneum (where most of the talc seemed to have accumulated) were studied by two methods. Sections were mounted, unstained, in Canada balsam and studied petrographically to determine mineralogical changes which had taken place. Also, sections were stained with hematoxylin-eosin and special stains to study histologically the changes in connective tissue.

From the mineralogical standpoint, two facts seem to warrant emphasis, according to the authors. First, most of the dolomite, which contained no silica, disappeared in 12 to 15 months with only the coarsest particles remaining. Second, in "talcs" which contained carbonate, the serpentine was also removed, but to a lesser extent. It was also noted that after a period of about 8 to 12 months talc needles tended to align themselves parallel to the abdominal wall of the animal.

Histological study showed that the material was mostly taken up and held by the giant cells. In all cases the fibrous tissue was of minimal amount; in no instance was there evidence of a progressive proliferation of the fibrous tissue similar to that found in the usual silicotic nodule. The pattern then is one of storage of foreign material, primarily in phagocytic cells, with sufficient accompanying fibrous tissue to hold these cells in place. The authors term this "storage fibrosis".

The two talcs which contained the least carbonate, 814-50 and 814-34 (see Table 8) caused the greatest amount of fibrous tissue. Studies of talcs which have been shown to cause granuloma will be reported in a later section.

C. Dogs

Newberne and Wilson (372), whose work on rats has already been reported, also treated purebred beagles with various silica compounds. The doses of each compound administered were equivalent to 0.8 g/kg/day of SiO_2 , mixed in a highly palatable diet. Table 9 shows the compounds administered, and the incidence of renal lesions

that were observed after sacrifice at four weeks.

Table 9 (372)

Renal Lesions in Dogs Fed Silicon Compounds for Four Weeks


Treatment	Dose (µm/kg/day)	Sex	Incidence of renal lesions
Control	0	M	0/6
Silicon dioxide	0.8	F	0/6
		M	0/9
Aluminum silicate	1.3	F	0/8
		M	0/6
Sodium silicate	2.4	F	0/7
		M	8/8
Magnesium trisilicate	1.8	F	7/8
		M	9/9
		F	9/9

The only significant clinical abnormalities exhibited by the animals during the treatment period were polydipsia and polyuria, seen in a few dogs fed sodium silicate and magnesium trisilicate. Soft feces, discolored by unabsorbed compounds, were seen occasionally in most of the test dogs. Body weight, food intake, and urinary and blood measurements were essentially normal in all animals. Upon autopsy, gross cortical lesions of the kidney were noted in all of the male dogs and in all but one of the female dogs that were fed sodium silicate. The lesions appeared to be focal, subcapsular hemorrhages, but on the cut surface they suggested cortical infarcts. Such lesions were seen in all dogs of both sexes that had received magnesium trisilicate, but were not observed in animals fed aluminum silicate or silicon dioxide. Histopathologic studies also revealed characteristic lesions from the sodium and magnesium compounds, but none in the animals receiving aluminum silicate or silicon dioxide. The nature of the lesion was the same

in all cases, although they varied in severity from one animal to another and from one area to another within a kidney. Selected tubules seemed to be affected and were often in juxtaposition to normal ones. Occasionally, deposits of crystalline material, characteristic of mineralization, were observed in degenerated tubular epithelium, but such changes were not common. Glomeruli did not appear to be damaged. The general impression gained by the authors was one of irritation of tubular epithelium, followed by degenerative and regenerative changes; these alterations were accompanied by inflammatory cell infiltration into the interstitium.

Finally, the authors emphasize that despite the presence of extensive renal damage, impairment of renal function was not detected by any of the chemical tests made on serum or urine. Apparently, renal reserve was adequate for normal functional demands. Perhaps longer exposure to the compounds, or examination after a longer period following the end of the treatment, would have shown impairment of renal function to be tied in with more advanced renal lesions(372).

D. Humans

The effect of  atmospheric concentrations of silicon dioxide on the health of children has been reported by Egorova (147). He studied the area around a factory manufacturing ferrosilicon of 45% and 75% concentration. The factory utilized open, electric-arc furnaces, with the furnace gases being discharged at the furnace tops; the gases consist mainly of the products of sublimation and vaporization of silicon and its compounds. Table 10 shows the dust concentrations in the atmosphere at distances from

its source, and also the free silica in the dust.

Table 10 (147)

Dust Concentrations in the Atmosphere at Different Distances from the Shop Producing Ferrosilicon Alloys

Distance from the source of atmospheric pollution, km	Number of samples	% of samples found to contain dust	Mean dust concentration found in the samples, mg/m ³	% of samples in which the dust concentration exceeded the max. permissible concentration of 0.5 mg/m ³
Shop grounds	23	100	4.1	
0.3	25	100	3.1	100
0.5	25	100	2.5	100
1.0	26	100	2.0	100
2.0	24	100	1.75	100
3.0	25	100	3.0	100

Concentration of Free Silica and its Percentage Contents in the Dust at Different Distances from the Shop Producing Ferrosilicon Alloys

Distance from the source of atmospheric pollution, km	Number of samples	% of samples found to contain free silica	Mean concentration of free silica in the dust, %	Concentration of free silica in the dust, %	
				mean	maximum
Shop grounds	23	100	0.72	14.0	41.6
0.3	25	100	0.54	16.0	36.0
0.5	25	88	0.30	11.2	31.5
1.0	24	83	0.37	20.0	66.0
2.0	24	87.5	0.19	11.8	32.0
3.0	25	92	0.50	14.7	45.0

To insure comparability of data, studies were made of the living and dwelling conditions of the children and the economic status of their families, as well as their location and the health patterns. The disease-incidence data for the children were based on requests for medical pediatric consultations, and the records of the medical consultation service. Comparative morbidity data showed that overall the children's morbidity was 37% higher in the investigated area near the factory than in the control area

during their first year of life, 27% higher during the their second year, 30% higher in the third year, 16% higher in the 4-7 year period, and 45% higher during the 8-15 year period. Ear, nose, and throat disorders were most common among the children residing in the test area. Data on child morbidity based on those regarding diseases are shown in Table 11.

Table 11 (147)

Morbidity in Children in the Investigated Area (incidence of diseases per 100 investigated children)

Statistical parameter	Overall morbidity	Disorders of respiratory organs	Otorhinolaryngologic disorders	Infectious diseases	Diseases of digestive organs	Other diseases
Control area						
$M \pm m$	107.5 \pm 0.63	15.35 \pm 0.80	37.7 \pm 1.00	46.9 \pm 1.10	4.29 \pm 0.45	3.98 \pm 0.43
Area with polluted atmosphere						
$M \pm m$	132.2 \pm 2.49 12	21.92 \pm 1.40 4.1	64.6 \pm 1.69 8.5	51.0 \pm 1.70 2.5	4.57 \pm 0.70 0.33	6.41 \pm 0.83 2.4

Regarding general development, the children in the tested area, showed a definite lag -- 71.8% of boys and 70% of girls showed medium and above-medium physical development in the tested area as compared to 91% and 89.6% respectively in the control district.

III. Long-Term Studies

A. Rabbits

Gardner and Cummings (176) designed experiments to demonstrate the relation of particle size to the amount of reaction produced by a measured quantity of silica. Quartz was administered to two groups of rabbits, eight animals per group, as 2% suspensions

(by wt) in physiological saline i.v. in the ear vein over a period of one to four months. The total dose for each animal was 1.3 g. Group I received silica particles 1 to 3 μ in diameter; Group II particles, 6 to 12 μ ; and Group III, aluminum oxide particles 1 to 3 μ in diameter. Seven of the 24 animals died of embolism during the course of treatment; two more in the group receiving the finer quartz particles died of silicosis. Aside from the one animal who died during the first injection from coccidiosis, the remaining rabbits were sacrificed two to three years after completion of treatment.

Histological examination showed that particles are segregated in different locations according to their size. The largest ones are caught in the pulmonary capillaries, those of intermediate size in the spleen and hepatic lymph node, and the finest ones in the liver. The authors indicate that the fine silica particles are the most active, producing progressive coarsely nodular cirrhosis of the liver, attended by extensive destruction of the parenchyma, and followed later by regeneration in certain areas. They indicate this cirrhosis is the result of a typical, hyaline nodular silicotic fibrosis originating in the portal connective tissues. The coarse particles, 10 to 12 μ in diameter, are much less irritating, exciting a simple foreign body reaction which progressed very little in three years. Fine aluminum oxide particles of the same size as those of the smaller silica particles merely are phagocytosed and produce no fibrosis in the stroma of any organ where they were deposited. The authors believe that

these observations indicate that the injury produced by silica is specific and chemical rather than physical in character.

B. Humans

Other than studies of silicosis as an industrial disease, long-term human studies on silica intake have been very rare. Akiya et al. (1966) have offered a detailed report on the relationship between silica in drinks and foods and in human blood vessels. As indicated in Section V, Chemical Information, silica is encountered by the human from much of his food and environment. The Japanese are excellent subjects for a study of this effect, because of their unusually high silica intake from talc milled rice, a staple of their diet, as well as from their drinking water. Akiya has proposed that, in addition to the calcium and cholesterol deposits, accumulation of silica in the blood vessels may be one cause of arteriosclerosis. He prepared and studied 89 aorta slides from Japanese patients. Table 12 traces the development of arteriosclerosis, with advancement in age. The author separated the aorta into three layers of the intima, media and adventitia, and determined the silica content of each layer. He noted that the silica content of the media seemed to remain somewhat constant, but that the silica content in the adventitia increased with advance of age, and its content in the intima decreased with advancing age. At the age of about 40 to 42 the quantities in these two layers reversed. It was presumed that the silica in the adventitia might have a connection with the hypertension or the arteriosclerosis that appear after 40 years of age.

Table 12 (006)

The Age, Sex, Cause of Death and View of Aorta of Samples

No.	Autopsy No.	Age	Sex	Cause of Death	View of Aorta
1	1329	3 Mon.	M	Erythroblastosis Fetalis	N.
2	1352	11 Mon.	F	Dystrophia	N.
3	1282	9	M	Acute Myeloid Leukemia	N.
4	1393	9	M	Subarachnoidal Hemorrhage	N.
(5)	16644	15	M	Poisoning of Hypnotics	N.
(6)	16724	16	M	Bacillary Dysentery	N.
(7)	16653	16	M	Drowning	N.
(8)	16671	17	M	Poisoning of Hypnotics	N.
9	1330	20	M	Postoperative Shock	N.
10	1323	20	M	Lung-metastasis of Osteosarcoma	N.
11	1391	20	M	Myeloid Leukemia	N.
12	1264	23	F	Rhabdomyosarcoma	N.
13	1280	24	F	Abdominal Tumor	N.
14	1366	24	M	Leukemia	N.
15	1324	25	M	Esophagus Stenosis	N.
(16)	16632	26	M	Poisoning of Cyanide	N.
17	1224	28	F	Rupture of Ectopic Pregnancy	N.
(18)	16643	28	M	Lobular Pneumonia	N.
19	1284	32	M	Sepsis	N.
20	1317	32	M	Purulent Meningitis	N.
21	1380	32	M	Hepatoma	N.
(22)	16717	34	M	Valvular Disease	N.
23	1342	35	M	Unknown	N.
24	972	35	M	Unknown	N.
25	1348	35	F	Gastric Cancer	N.
26	1370	37	F	Pulmonary Cancer	N.
27	1274	38	M	Neuroblastoma	N.
(28)	16633	38	M	Acute Heart Disease	N.
(29)	16683	38	M	Fracture of Skull	N.
(30)	16694	40	M	Unknown	A(+)
(31)	16696	40	M	Cerebral Bleeding	A(+)
(32)	16629	41	F	Poisoning of Hypnotics	N.
(33)	16624	41	F	Rupture of Tubal Pregnancy	N.
34	378	42	M	Teleangioma of Femoral Bone	N.
35	1260	42	M	Pulmonary Cancer	A(+)
36	1340	42	F	Gastric Cancer	N.
37	1332	43	F	Liposarcoma	N.
(38)	16697	43	M	Acute Cardiac Insufficiency	N.
39	1347	44	M	Aplastic Anemia	N.
40	1296	46	M	Cancer of Maxilla	N.
41	1364	46	M	Pancreatic Cancer	A(+)
42	1346	47	F	Ovarial Cancer	N.
(43)	16710	48	M	Acute Heart Disease	N.
44	1261	49	M	Aortic Aneurysm	A(++) Ca(++)
45	1265	49	F	Ovarial Cancer	N.
46	1257	49	M	Gastric Cancer	A(++)

Table 12 (Cont) (006)

No.	Autopsy No.	Age	Sex	Cause of Death	View of Aorta
17	1258	49	M	Cerebral Bleeding	A(II)
48	1383	49	M	Gastric Cancer, Pulmonary Tuberculosis	A(+) Ca(+)
49	1168	50	F	Uterine Cancer	A(III) Ca(III)
50	1278	50	M	Gastric Cancer	N.
51	1263	51	M	Esophageal Varix	A(III)
52	1361	51	M	Auricular Fibrillation	N.
53	1267	51	F	Hepatic Cancer	A(II)
54	1266	52	M	Mesothelioma	N.
55	1262	53	M	Pulmonary Cancer	A(II) Ca(+)
56	1152	53	M	Liver Cirrhosis with Hepatoma	A(III) Ca(+)
57	1281	54	F	Arrhythmia	A(III) Ca(III) U.
58	1336	54	M	Pulmonary Cancer	A(II)
59	1313	55	F	Hepatic Cancer	A(II)
60	1304	55	M	Mucocpidermoid Cancer	A(III) Ca(III) U.
(61)	16652	55	F	Acute Hemorrhagic Bronchitis	A(II) Ca(+)
62	1212	56	M	Wilson's Disease	A(II) Ca(III) U.
63	1285	56	M	Miliary Tuberculosis	N.
(64)	16686	56	M	Coronary Sclerosis	A(II) Ca(+)
65	1234	57	M	Gastric Ulcer	A(III) Ca(III) U.
66	1326	57	F	Gastric Cancer, Renal Tuberculosis	N.
67	1315	58	M	Pulmonary Tuberculosis	A(III) Ca(III) U.
68	1374	58	M	Cancer of Mandible	A(II)
69	1290	60	M	Cancer of Maxilla	A(II) Ca(II)
70	1289	60	F	Cancer of Rectum	A(II)
71	1306	62	M	Pulmonary Tuberculosis	A(II)
72	1277	62	M	Unknown	A(+)
73	1358	62	M	Cancer of Colon Transversum	A(III) Ca(III) U.
74	1308	62	F	Myocardial Infarction	A(III) Ca(III) U.
75	1269	63	M	Prostatic Carcinoma	A(III) Ca(III) U.
(76)	16658	63	M	Acute Cardiac Insufficiency	A(III) Ca(III) U.
77	1104	64	M	Pulmonary Cancer	A(II)
78	1279	64	F	Abdominal Tumor	A(II) Ca(II)
79	1379	64	M	Death after Lobotomy	A(+) Ca(+)
80	1389	65	M	Cancer of Urinary Bladder	A(III) Ca(III) U.
(81)	16685	66	F	Acute Coronary Insufficiency	A(II) Ca(+)
82	1310	67	M	Pulmonary Cancer	N.
(83)	16702	67	F	Acute Pneumonia	A(II)
84	1355	68	F	Cancer of Colon Descendens	A(+)
85	1372	69	F	Hepatic Cancer	A(III) Ca(III) U.
86	1381	69	M	Gastric Cancer	A(III) Ca(III) U.
87	1388	70	F	Paralysis of Medullaoblongata	A(II)
(88)	16700	74	M	Drowning	A(III) Ca(III) U.
(89)	16703	74	F	Lobular Pneumonia	A(II)

Aortae supplied by Tokyo Medical Examiners Office are indicated No. in (). Others were supplied by Department of Pathology, Tokyo Medical and Dental University.

View of Aorta: N....Normal, A....Atherosclerosis, Ca....Calcification, U....Ulcer
(+)...slight degree, (II)...middle degree, (III)...high degree.

In general the authors also found that the human organs in the cases of arteriosclerosis contained larger amounts of silica than in other cases. Other results from this study are discussed in later sections of this monograph.

Silicosis as an industrial disease has long been recognized; in view of the great bulk of literature on the subject, and the general knowledge concerning it there seems no need to consider that condition, when it results from industrial exposure, in this monograph. When certain aspects of the disease shed light on the effects of silica from other types of exposure, there will be discussion of it.

IV. Special Studies

A. Carcinogenicity

It is well established that while cancer of the stomach in man is one of the most common of all tumors, in animals it is rare (475). Also, within the animal group, rats have been used almost exclusively for study since guinea pigs, mice, and rabbits do not develop gastric tumors. The lesions in the rat forestomach and glandular stomach usually have been described as papilloma presumably because the relatively short lifespan of the animals does not give sufficient time for malignant development.

The work of Sugiura (475), when considered in the above context, offers confirmation of the possible carcinogenic stimulation of a silicon mixture. Young albino rats, Sherman stock, from 30 to 75 days old and weighing from 60 to 125 g were placed on a mixture of 95% polished rice and 5% kieselguhr (infusorial or diatomaceous earth). This dietary regime was maintained for their lifespan. The results of this study are summarized in Table 13, and illustrate clearly that the daily oral administration of kieselguhr has a definite stimulating effect on the production of stomach lesions.

Table 13 (475)

**Summary of Incidence of Lesions Observed in the Forestomach
of Rats Fed Rice Alone or Rice and Kieselguhr**

Experiment No.	Diet	Number of rats used	Age of rats, days	Duration of feeding, days	Number of animals with lesions	Number of animals without lesions
1	White rice and kieselguhr	10	30	49-91	9	1
2	White rice and kieselguhr	11	75	98-116	11	0
3	White rice	27	30	54-148	13	14
4	White rice	24	75	65-156	16	8
5	Brown rice	12	30	54-75	0	12
6	Brown rice and carrot	30	75	88-200	0	30

Attention is called to the negative results of Bertke [(061) Short Term Studies section, 1] on Wistar rats fed diatomaceous earth 5% in their diet. No abnormal lessions were found in the stomach, small or large intestines after 90 days of treatment and then immediate sacrifice, possibly a result of the difference in strain.

As mentioned earlier in this monograph, the Japanese prefer talc milled rice (342), and this fact has been tied to the very high incidence of stomach cancer in Japanese men. Merliss (342) quotes from work by Hirohata and Kuratsune (Brit. J. Cancer 23, 465, 1969) a rate of 67.96% per 100,000 in the male population of Japan, 7 times that of the rate for men in the United States, and the highest of the 24 nations studied.

There are numerous reports in the literature of granulomas, at times fatal, from talc used in surgical procedures (as surgical glove powder). Ross and Lubitz (421) have published a review of its incidence up to 1949. For the purposes of the monograph, a single case of talc granulomata attributed to earlier surgery should suffice to illustrate the point. Diffenbaugh (139) has reported the case of a 59-year-old woman admitted to the hospital with a history of nausea, vomiting and a weight loss of 60 lbs in one and a half years. She had a history of a pelvic laparotomy eleven years earlier at which time adhesions from about the gall-bladder were freed. Exploratory laparotomy at the time of the second admittance revealed numerous small and large grey nodular masses covering the entire peritoneal surfaces and omentum, with dense diffuse adhesions throughout the abdomen. There was no ascitic

fluid present and the stomach was normal. The nodules were removed, and her post-operative course was uncomplicated, with no recurrence of nausea or vomiting and with a pattern of weight gain. Histological examination of sections of the nodules showed a large amount of fibrillar scar tissue, with many large and small vacuoles at the margins of which were foreign body giant cells. Scattered in the dense fibrous portions were other small clusters of foreign body giant cells and large mononuclear phagocytes. Some of the giant cells had fragments of refractile material which, examined under the micropolariscope, showed a needle-like structure with double refractive properties. Another significant aspect was a mass of concentrically arranged hyalinized fibrous tissue with small crevices in the center.

The diagnosis was that of extensive foreign body granuloma, and the author postulated that physio-chemical changes in the tissues cause increase in soluble fractions of the talc which stimulates the tissues to reaction. He pointed out that similar delay of pathological changes in silicosis of the lungs is attributed to slow leaching of the silica.

A case of metastatic carcinoid tumor associated with complicated silicosis has been reported by Halonen et al. (213). The patient was a 53-year-old widow who had worked in a china factory for some six years, starting at age 28. At age 31 she had lost weight, suffered from fatigue etc., and a diagnosis of goiter was made (a condition also attributed to silica intake (493)). She was operated on at that time and the symptoms disappeared. At age

48, she suffered from acute tonsillitis and at 49 a routine chest X-ray revealed a shadow in the right upper lobe. Edema of the legs, flushing and then painful nodules of the legs developed shortly before her admission to the hospital at age 52. X-ray examinations at this time indicated tumor-like shadows in both lungs and a tumor projecting into the lumen of the cecum. A laparotomy was undertaken, and a hard tumor discovered. It was judged inoperable because when it was manipulated a severe bronchoconstriction and convulsive contractions of the small bowel suddenly occurred. The blood pressure fell to shock level, but was restored by the administration of adrenaline chloride. Samples were taken from the metastases for microscopic study; and these subsequently showed the features of a carcinoid tumor. She recovered sufficiently to return home, but returned in 5 months and died two months later. The primary tumor proved to be at the end of the ileum with metastatic nodules in the liver, spleen, mesentery, omentum, both ovaries and in the Douglas pouch. The lungs were fibrotic throughout and contained coal pigment, with double refracting particles seen under polarized light. The authors point out that the development of silicosis usually takes exposure to silica dust for at least 10 to 15 years. It is known that silicosis may progress after exposure ends, with infection usually held responsible. Many of those cases are considered to be tuberculous. In summary, while the patient's exposure to silica dust had been of only six years duration, and it ended some 15 years before the tumor-like shadow appeared in her lung, the autopsy

showed she was suffering not only from a metastatic carcinoid tumor but from complicated silicosis which had considerably progressed several years after exposure to silica had ended. The authors suggest the possibility of the action of 5-hydroxytryptamine on the development of the silicosis.

As already indicated, surgical glove talc granulomas in various intestinal and pelvic organs have been reported. Localized silica granulomas due to glove talc also have been observed in surgical scars and as a result of impregnation of the skin by siliceous particles such as glass or sand (496). However, the report of extensive granulomas of the skin seem to be somewhat unusual, and Tye (496) described the case of a 45-year old woman, admitted to the hospital because of the development of increasing numbers of lumps on her shoulders, buttocks and thighs. The nodules had been developing for about two years and were at sites where she had had as many as 500 boils over a span of 8 to 10 years. The boils had been treated by incision and drainage, and it was observed that the nodules did not appear on areas where there had been no furuncles. It had been her practice for several years to take one or two baths daily, followed by liberal dusting of her entire body with powdered talc. Hundreds of hard, slightly raised, flesh-colored nodules varying in size from 2 mm to 2 cm were scattered over her back, buttocks and anterior and posterior aspects of her thighs. Histologically, the lesions were composed of islands of granulomas with numerous foreign body cells. From electron microscope and x-ray diffraction studies it was concluded that the

foreign bodies in the phagocytic cells were talc. It was postulated that the talcum powder gained entrance through the skin at the sites of the draining or incised boils.

B. Teratogenic Effects

Ludox[®], a suspension of colloidal silica from the Du Pont Co. which contains 15% SiO_2 , 1% Na_2O , 0.001% NaCl and 0.003% Na_2SO_4 , has been tested for teratogenic effects in the amniotic cavity of chick embryos (526). The silica particles were dense, nonporous spheres approximately 7 μ in diameter. They remained discrete and maintained particle size on dilution in water or saline. The pH was 8.5 with minimum stability range at pH 5-6. Fertile eggs from White Leghorn flocks or from Meatline hens mated to New Hampshire roosters were incubated at 37.5°C, and the Ludox solution was injected into the amniotic cavity of eggs incubated for five days. Each egg received 0.05 mg of the diluent containing calculated amounts ranging from 0.001 to 0.25 mg. Controls, with the same incubation setting and period, were treated with saline or distilled water with the pH adjusted to that of the test material. The embryos were observed through the shell openings and dead embryos removed for gross examination. Surviving embryos were sacrificed after 15 days, and incidence and description of defects were tabulated for all embryos surviving to the 8th day of incubation. Table 14 indicates the survival and percent of abnormalities resulting from the Ludox inoculation.

Table 14

**Survival and Incidence of Abnormal Development in Chick Embryos
Inoculated with Colloidal Silica into the Amniotic Cavity at
5 Days Incubation (526)**

mg contained in 0.05 ml diluent per egg	Total eggs	Embryos surviving to incubation day:		Total abnormal of those surviving 8 days or more.	
		8	15	No.	(%)
Carbon 0.1	73	38	10	19	(50)
" 0.01	52	30	13	5	(17)
" 0.001	51	20	9	1	(3)
Control	72	51	27	0	

The defects included retarded lid development or exophthalmia, encephalocoele or exencephaly, external edematous blebs, ectopic viscera, retarded feathering and crossed or short upper beak. Also, there were severe axial distortions and correlated limb distortions that the authors emphasize were similar to those that have been observed with non-toxic colloidal alumina.

C. Effects on Action of Bacilli

The relationship of silicosis and tuberculosis has been considered by a number of authors, as already mentioned. Detailed discussion of their findings seems outside the scope of this monograph. However, in view of the fact that silica can enter the body from the environment either through the lungs or the digestive tract in normal daily living, and circulates via the blood stream into tissues as well as being excreted via urine and feces, this relationship could be a factor in the overall picture of toxicity. For that reason the studies of Cummins and Weatherall (120) are mentioned. They have reported that in vitro studies with tubercle bacillus in both saline and silica sol cultures indicated that the growth of tubercle bacilli in blood in vitro is neither favored nor

inhibited by the presence of colloidal silica. This conclusion seemed to be in some disagreement with the earlier work of Gye and Kettle [Brit. J. Exp. Path. 3:244 (1922)] who reported that tubercle bacilli injected into the circulation of rabbits and mice tended to settle in localized subcutaneous lesions produced by various agents which brought about increased vascularity and tissue necrosis. The latter authors further concluded that the destructive action of silica on cells, whereby the foci of necrosis was produced in which tubercle bacilli could multiply, was actually a case of cell poison by a soluble derivative of silica.

In regard to other bacilli, Cummins and Weatherall (119) studied the role of colloidal silica in bactericidal action against typhoid bacilli. They concluded from their studies that colloidal silica solution preserves typhoid bacilli from lysis in human blood or serum, and that this property extends to other serum-soluble bacteria. In their experiments they found colloidal silica was able to preserve Type I pneumococci from lysis in ox bile. Also, they reported that the addition of colloidal silica solution to a cellular exudate, produced by the intraperitoneal injection of dust particles, leads to agglutination of the leucocytes and dustcells and to the preservation of a large proportion of these cells, intact though probably dead, for a period of two months or more. Finally, they concluded that it is probable that these anti-lytic and agglutinating properties of colloidal silica solution may, through interference with cytolysis, be factors in the production of the lymphatic obstruction associated with pulmonary silicosis in man.

BIOCHEMICAL ASPECTS

I. Breakdown

No information on spontaneous changes in the silica compounds or mixtures in food storage or processing was available from the survey made for material for this monograph. Regarding changes in the tissues of living animals, reference is made to the work of Schulz and Williams (44) discussed in the Biological Data section, Short Term studies. These authors reported the petrographic changes which took place in seven different commercial talcs injected peritoneally in guinea pigs. They observed that the amount of talc itself (magnesium silicate) was not disturbed in the 15-month span of the study, but that the other substances in the commercial samples, notably dolomite and also serpentine when in the presence of carbonate, were slowly leached out of the particles as they remained in the animal tissue.

II. Absorption and Distribution

A. Rats

1. Kochmann and Maier, utilizing the rats followed for growth on the four different forms of colloidal silica [(273) discussed in the section on Short Term studies] studied the whole body retention of silicic acid. Four or five animals of each series were sacrificed, ground in a meat grinder, the fresh substance weighed, dried over a water bath, and rubbed to a fine powder. The water content was determined, as well as that of fat, and nitrogen, with the silicic acid being determined by the method of Schulz [Pflugers Arch. f. d. ges. Phys. 144: 350 (1912)]. The results of these analyses are shown in Table 15.

Table 15 (273)

Study on Whole Body Retention of Silicic Acid in Rats

1 Fütterungsart der Ratten	8 Naß- substanz g	9 Trocken- substanz in 100 g Naß- substanz	10 Fett in 100 g Naß- substanz	11 Wasser in 100 g Naß- substanz	13 In 100 g reiner Trockensubstanz sind enthalten										18 In 100 g Asche SiO ₂ %
					12 Asche		14 Asche Mittel- wert %	N ₂ %	15 N ₂ Mittel- wert %	16 Eiweiß Mittel- wert %	SiO ₂ I II %		17 SiO ₂ Mittel- wert %		
					I. %	II. %					I. %	II. %	I. %	II. %	
2 Silvana, 5 Ratten	850,00	28,40	4,20	72,20	15,52	15,65	15,58	I. 14,95 II. 15,65	15,30	95,63	I. 0,015 II. 0,012	0,014	84	A	
3 Kontrolle zu Silvana, 4 Ratten	705,00	21,20	6,60	72,05	17,76	17,89	17,83	I. 13,12 II. 13,12	13,12	82,09	I. 0,0123 II. 0,0167	0,0145	79		
4 Silistren, 4 Ratten	498,00	20,98	9,61	69,40	16,68	16,79	16,74	I. 15,01 II. 15,01	15,01	93,81	I. 0,0379 II. 0,0431	0,041	259		
5 Siliquid, 5 Ratten	694,00	22,58	9,54	67,94	14,11	15,65	14,88	I. 13,74 II. 13,74	13,74	85,87	I. 0,0204 II. 0,0199	0,0201	196	B	
6 Kolloidale Kiesel- säure, 5 Ratten	676,00	21,04	8,91	70,04	15,49	16,54	16,01	I. 14,66 II. 14,66	14,66	91,63	I. 0,0293 II. 0,0273	0,0286	179		
7 Kontrolle zu Sili- stren, Siliquid u. kolloidal. Kiesel- säure, 5 Ratten	671,00	20,23	8,58	71,78	16,78	16,03	16,41	I. 14,78 II. 15,10	14,94	93,37	I. 0,027 II. 0,023	0,025	144		

1. Type of feeding which rats received
2. Five rats received water from Silvana mineral springs
3. Four rats received water from Silvana mineral springs-Control
4. Four rats received Silistren (trade name)
5. Five rats received Siliquid (trade name)
6. Five rats received colloidal silicic acid
7. Five rats received Silistren, Siliquid, and colloidal silicic acid-Control of
8. Wet substance
9. Dry substance in 100 g wet substance
10. Fat in 100 g wet substance
11. Water in 100 g wet substance
12. Ash
13. Contained in 100 g dry substnace
14. Average wt. of ash
15. Average wt. N₂
16. Average wt. protein
17. Average wt. SiO₂
18. In 100 g SiO₂ ash

The authors pointed out that there was no appreciable difference between individual animals in their water content; that the fat content of older animals (series 1) was lower than of the younger ones (series 2); and that there were no characteristic differences between the groups of animals in fat-free substances, nitrogen content, or generally, ash content. They did emphasize however, that the silicic acid content of the dry substance and the ash content was lower in the heavier older animals than in the younger lighter ones, a finding which agrees with that of Schulz (originator of the analytical method for silicic acid) who always found larger quantities of SiO_2 in the organs of younger subjects. There was no apparent retention of SiO_2 in the animals fed this in the form of natural mineral water, but those fed with Silistren, and possibly those fed with colloidal silicic acid, showed SiO_2 accumulation. They were puzzled as to why Siliquid-fed animals apparently retained none. Finally, they stressed as worthy of note that such small quantities of SiO_2 , the regime of their experiments, could result in any accumulation.

2. Webb, Salle and Tienes (511) in their study of rats fed SiO_2 as Braun's Air Float, 325 mesh, in their food, found some silica accumulation in liver, spleen and testis. The procedure of King and Stantial [Biochem. J. 27: 990 (1933)], with modification of pH control to insure complete removal of phosphate, was used; the results are shown in Table 16.

Table 16 (511)

Silica Content of Tissues

ORGAN	SERIES I	SERIES II	SERIES III	SERIES IV
Liver.....	19.0*	22.7	25.0	40.5
Spleen.....	10.4	16.8	42.5	59.5
Testis.....	17.7	18.1	51.8	47.3
Kidney.....	22.7	16.3	25.0	22.3
Lung.....	34.1	22.2	37.0	25.9
Eye.....	64.2	49.6	58.8	57.1
Blood.....	14.1	17.8	13.3	15.1
Average.....	19.8	20.6	32.4	35.1

* Mgm. SiO₂ per 100 gm. of fresh tissue.

These authors interpreted their results on human tissues as being in general accord with the results obtained by others, as summarized in the review article of Belt and King (262). Also, they point out that since silica is generally assumed to circulate in the blood in colloidal form it is not surprising that it is picked up predominantly by organs containing a large amount of reticulo-endothelium tissue. In regard to the eye, they note that while it contains the highest concentration of silica of any organ investigated, this seems to be a normal constituent, with added silica not accumulating there. By a series of calculations Webb et al. (511) proposed the fraction of the ingested silica that is absorbed through the gastro-intestinal tract and accumulated in the tissues. When silica was added to the food to the extent of 0.1% the quantity accumulated by the tissues represented 0.8% of the amount ingested. In series III, silica analyses were made on urine and feces for one week, and on the basis of these results, the authors propose that, of the added ingested silica, 95% was excreted in the feces, 4% in the urine, and 1%

accumulated in the tissues.

3. The lung as a portal of entry into the circulatory system has been illustrated by the study of Byers and Gage (084). Experimental details have already been given in the Short Term Studies section, 5. The three samples tested, A1, particle size 19 μ , A2 particle size 20 μ and Sample B particle size 25 μ were administered intratracheally and the rats sacrificed at intervals up to 1 year. Silica was determined in various organs by modification of the method of King, Stacy, Holt, Yates and Pickles (257) described in the Analytical Methods section of this monograph. Accumulation data is reported in Table 17 below:

Table 17 (257)

Retention of Silica by Rat Tissues

Sample	Weeks After Injection	Lungs	Liver	Kidneys	Spleen
A1	12	1,570	177	33	0
	24	755	20	12.4	1.4
	52	210	61	9.7	0
A2	12	5,150	249	138	0
	36	1,550	153	44	19
	52	720	153	56	0
B	12	652	234	34	5.8
	24	324	43	8.9	1.4
	52	108	38	12	0
Average silica content of normal rat tissue		28	37	11	5

All rats received an intratracheal dose of 25 mg. precipitated silica. Figures represent μ g. SiO_2 per rat.

As can be seen from the data, silica disappeared progressively from the lungs, although it was still found in significant amounts after 12 months. Some initially appeared in the liver and kidneys of animals in all three groups, but only in A2 was silica noted in these organs after 6 months. The authors say they find no evidence

to indicate the route by which silica is eliminated from the lungs, and whether it is transported in the body as particles or in solution. It would seem appropriate, however, to call attention to the fact that A2, with its greater tendency to aggregate [as discussed in Short Term Studies Section ^{A5}] had the most marked effect on lesion formation, and the proportion retained by lungs, liver and kidneys was the greatest for the three samples.

B. Rabbits

1. Intravenous administration of silica has been studied by Mosinger, Jouglard-Duplay, Versino and Granier (359). Crystalline silica was graded by sedimentation in water, dried, sterilized, and checked under the microscope. The authors indicate the particle size to be in the range from 1 to 5 μ , with ca 80% 3 μ particles. It was suspended in "physiological solution" and injected i.v. into the marginal ear vein of tawny Bourgogne strain rabbits. The quantity administered per injection was 100 mg, with the total dose being 500 to 600 mg spaced over a period of 10 to 15 months. The animals were 3 months old at treatment initiation, and weighed 1,200 to 1,500 g (with a normal weight development noted during the experimental period). Silica in the blood was measured by a modification of the procedure of King [Bull. Soc. Chem. Biol. 12:903-909, 1939 and Biochem. J., 56:11-16, (1954)]. Normal values for untreated rabbits were measured, varying from 0.305 to 0.770 mg/100 ml blood. This averages 0.454 mg/100 ml as contrasted with the high normal of 0.400 for man [Revue Lyonnaise de Medecine, 8:1375, (1959)]. This difference the authors attribute to the

exclusively vegetable diet of rabbits. In the treated rabbits, 9 of 13 showed silica of 1 g/100 ml in the blood; when silica content was plotted as a function of time elapsed since the last injection, a decreasing curve was obtained which showed that beyond a period of two months after the last treatment the silica content of the blood reverted to normal in a majority of the cases. These authors then conclude that it is reasonable to suppose that the silica injected i.v. is localized in the receptor organs. Autopsies of the rabbits, were made and the procedure of Isaacs [Bull. Soc. Chim. Biol. 6: 157-168 (1924)] used to determine silica in specific organs. For the liver, a range of 8.8 mg to 435 mg/100 g dry tissue was found, and for the lungs 2.6 mg to 278 mg; these values emphasize the mechanical stoppage of the silica particles in the pulmonary capillaries and also the process of liver detoxication. The spleen was also heavily loaded, with values up to 200 mg. Smaller quantities were found in the kidneys, with a maximum accumulation of 37.7 mg in the parenchyma. Protein disturbances and hormonal changes, as well as lesions of the spleen, lymph nodes, lungs and aorta were noted and are discussed in a later section.

C. Humans

1. King, Stantial and Dolan (259) have reported on the silica content of tissues from humans from various age groups, and also from individuals with a variety of health conditions. Fetal tissues have shown the following silica values [as determined by the procedure of King and Stantial, Biochem. J. 27: 990 (1933)].

Table 18 (259)

Silica Content of Fetal Tissues

Tissue	MgSiO ₂ /100 g dry tissue
Human: lung	9.8, 8.4
liver	3.8, 5.6
kidney	13.3
spleen	20.3
heart	20.4
brain	22.0

The normal paths of entry for silica into the body are through the lungs and the digestive tract; the greater portion of that entering by the latter route is eliminated in the feces. Some enters into the bloodstream; given the state of knowledge at that date, these authors believe this fraction was excreted in the urine without much retention by any organs of the body. However, a number of other workers have shown specific data on retention which, at times, is considerable (084,511,273 and others). As with the reports on the analysis of any substance, particularly in biological material, the analytical method is a critical factor; Table 19 (359) illustrates this statement. [REDACTED] Normal values for silica in rabbit blood by four different methods are shown in Table 19.

Table 19 (359)

Determination in the Rabbit
(in mg silica per kg or liter of tissue or fresh blood)

	Ramel	Isaacs	King	King et al. Stantial
Brain	0	8.8 to 18	—	—
Blood	12	—	—	3.4 to 5.8
Liver	35	0 to 5.6	150-212	96-57
Kidney	28	5-17.5	115-157	67-76
Lungs	270	3.9-12	125-139	85

Also, particularly in the case of entry through the lungs, the form of the silica, (the nature of its chemical compound, particle size etc.) are factors which can cause apparent discrepancies. This point is discussed later.

In spite of the uncertainty described above, it is of value to report the silica content of human and other animal tissues, as published by King, Stantial, and Dolan (259). Table 20 gives this information

Table 20

Silica Content of Tissues (259)

Tissue	mg. SiO ₂ per 100 g. dry tissue	Tissue	No.	mg. SiO ₂ per 100 g. dry tissue
Human:		Rabbit:		
Lung* (normal adult)	140	Lung	430	148, 101
(pneumonia)	93		360	235, 229
(silicotic)	3640	(young)	378	18.4, 14.2
"	1700	(animal exposed to SiO ₂ dust)	651	6755, 7115
"	880	"	693	805, 785
"	1630	"	687	1002
"	810	Liver	687	12.4
Liver* (stone grinder)	196	"	648	30.6
Spleen*		"	378	21.4
(normal)	15.1	"	651	20.8, 14.3
(septicæmia ♂ 30)	16.0	"	360	20.0
(bronchopneumonia ♀ 75)	43.0	Kidney	373	16.4
(rheumatic heart ♀ 46)	14.2	"	380	22.2, 17.8
(bronchopneumonia ♀ 36)	23.0	"	647	14.4
(pulmonary embolism ♀ 67)	30.0	(animal exposed to SiO ₂ dust)	651	50.0, 47.6
(carcinoma ♂ 25)	45	Lymph glands		80
(silicosis, fibrosis of spleen ♂)	345	Spleen	657	16.0
(silicosis, stone grinder)	436	(animals exposed to SiO ₂ dust)	360	22.5
Mesenteric lymph gland*		"	369	76.0
(stone grinder)	1080	Hair (grey)		16.5
Bone-marrow	26.0	(white)		15.6
(lumbar vertebra)	25.2	Guinea-pig:		
Ox:		Lungs		78.0
Lung	103, 100	"		48.0
"	104, 120	Hair		19.4
"	87.2, 81.2	"		20.7
Liver	21.6, 18.0	Chicken feathers		15.1
"	12.0, 15.0	"		12.9
Dog:				
Lung	115			
"	270			
Kidneys	28.7			
"	31.8			

* These analyses were performed on petrol- and ether-extracted tissues obtained from the wax blocks in which they had been imbedded after fixation for microscopic section and do not represent the silica content of the tissue as a whole. We are indebted to the Department of Pathology for this material.

2. King and Belt (256) have further studied the accumulation of silica in the lungs of 35 persons employed in dusty occupations and in the peribronchial lymph glands of 132 cases of individuals having no history of exposure to industrial dust. In some cases the concentration of silica was as high in lungs with no silicotic nodulation as in those with silicotic lesions. They found no evidence of a direct association between concentration of silica and the degree of fibrosis; they also indicate that the heavy deposits of siliceous particles found in the glands of old people have not produced significant silicotic lesions.

3. An extreme case of magnesium and aluminum silicate widely disseminated throughout a human body was that of a drug addict with a history of protracted use i.v. and s.c. of paregoric in various forms, combinations, and concentrations (obtained from proprietary drugs) (083). Paregoric, in proprietary preparations such as Donnagel and Parepectolin, when crushed and injected can cause the user to receive talc (hydrous magnesium silicate) and kaolin (aluminum silicate). The man died eight hours after entry into the hospital with a typical clinical picture of tetanus, a common occurrence in s.c. drug addicts. Upon autopsy, skin sections showed numerous foreign body granulomas, with large birefringent crystals. The lungs showed foreign body granulomata within the pulmonary arterioles. In the larger pulmonary arteries angiomatoids were present. Birefringent foreign body materials were in the walls of the development as well as within the walls of the venous channels in positions adjacent to the lumina. X-ray diffraction of the material

identified it as magnesium silicate. In the liver there was extensive periportal fibrosis, with numerous pigmented macrophages containing double refractive crystals; however, no foreign-body giant cells were seen. Foreign-body material was enclosed in the macrophages of the spleen. Also, the central nervous system disclosed extremely fine, double refractile particles in nerve bundles entering the anterior roots in the cervical region. No reaction was present in these fibers. The author indicated he did not understand the absence of foreign body giant cells in the liver, and postulated that perhaps the fine particulate size of the silicate crystals, which pass through the filter of the lungs into the peripheral circulation, favors engulfment by macrophages, while the larger crystals remaining in the lungs generate the foreign body reaction.

III. Metabolism and Excretion

At present, metabolism of the various compounds of silicon that are candidates for human intake is for the most part not a definable process. The exact fate of a mineral salt such as magnesium trisilicate has been predicted by Kas'yanenko (249). He has proposed that magnesium trisilicate becomes "jelly-like" upon entering the stomach as a result of the formation of colloidal silicic acid from the neutralization of hydrochloric acid. Page, Heffner and Frey (380) further postulated that when a silicate is acted on by an acid, part of the silica so formed is precipitated as a gel and part remains in solution as a colloid. It is then probable, they state, that as a silicate passes through the digestive tract other breakdown

products are formed such as ortho-silicic acid (H_4SiO_4) which is soluble, meta-silicic acid (H_2SiO_3) and tri-silicic acid ($\text{H}_2\text{Si}_3\text{O}_8$) which are partially soluble, and di-silicic acid ($\text{H}_2\text{Si}_2\text{O}_5$) which is practically insoluble. Numerous ideas and theories of what takes place with some of the other substances in vivo have been reported in the literature and are discussed in a later section of this monograph. Excretion in a more or less straightforward context, with reference to Si, is discussed here.

A. Rats

1. Holt, Yates and Tomlin (231) utilized ^{31}Si to follow the excretion of silicic acid in rats. They found there was rapid absorption of ^{31}Si from the peritoneal cavity, and analyses of separate tissues indicated no evidence of accumulation in any of the organs other than the kidney. They postulated that the concentration in the kidney was due to the tubular re-absorption of water and the non-absorption of silicate from the glomerular filtrate. They supported this theory with autoradiographs of the kidney slices which showed general distribution of ^{31}Si throughout the organ but with a higher concentration in the medullary portion where there were tubules only. When they used the liver concentration as representative of the concentration in other parts of the body, the time-concentration curve for the kidney after the injection of 8 ml of a solution containing 1 mg SiO_2/ml showed a sharp rise to the 3 hour value, followed by a gradual decline. The liver merely showed a steady drop in concentration down to a value comparable with that found in normal tissue. Varying amounts of silicate gave a

similar sharp rise with increasing dose, followed by a rapid fall to a value even below that for liver. From this they concluded that at low concentrations the silicate was concentrated by the kidney and rapidly excreted. When injections were given at higher concentrations, the body apparently was unable to deal with it, probably because the silicate is highly polymerized.

2. The effect of Aerosil, when injected i.p. into rats, was studied by Arnold, Sasse and Strecker (1941). 1000 mg in a 1% suspension was administered, with sacrifice at various periods, and sections for histographic study were prepared of the collecting tubule, glomerulus, proximal tubule, Henle's loop, and the distal tubule. The first series of studies was made after injection of Aerosil of surface area $150 \text{ m}^2/\text{g}$. The experiment was repeated with Aerosil O x 50 ($38 \text{ m}^2/\text{g}$) (about four times coarser than the first Aerosil) and also with silicic acid gel. Upon studying the sections of the three different series of experiments, the authors were surprised to find that in the lumina of the collecting tubules, particles of the same shape and dimensions were found in both the Aerosil groups; also, particles appeared in the collecting tubules of the silicic acid gel group which could not be differentiated from the particles from the Aerosil injections. The authors interpreted this to mean that from the surface of the Aerosil silicic acid enters into solution that polymerizes in the distal nephron sections, and they further postulated a glomerular excretion of silicic acid from all three initial sources.

B. Guinea Pigs

1. Four different forms of silicates were administered to guinea pigs by Sauer, Laughland, and Davidson (433), and the urinary and fecal excretions were measured.

a. Sodium metasilicate ($\text{Na}_2\text{SiO}_3 \cdot 5\text{H}_2\text{O}$) in a single dose corresponding to 80.0 mg of SiO_2 was administered orally; also four doses of the sample, each 80.0 mg of SiO_2 were given at 48 hour intervals. Silica in urine and feces was determined by the procedure of King et al. (257 of this monograph), and the "net silica excretion" (the difference between silica ingested and the silica excreted in the urine and feces) was calculated. For the single dose, given to a group of five pigs, the urinary excretion rose to a maximum of 9.7 ± 0.86 mg 48 hours after dosing, and returned to normal levels after eight days. The net silica excretion amounted to 48.3 ± 3.58 mg SiO_2 and represented ca 60% of the administered dose. After the administration of the first of four successive 80-mg doses the urinary silica excretion reached a maximum of 10.3 mg per 48 hour period, and did not increase with subsequent doses. A leveling off trend began 48 hours after the last dose, and silica excretion approached normal after eight 48-hour periods. At that time the net silica excretion was 307.0 ± 7.85 mg or ca 96% of the total dose.

b. A 2% silica sol, prepared from a solution of sodium silicate and dilute HCl, was administered i.p. in two dosages (100 mg and 60 mg SiO_2) to three and two pigs respectively. It was found

that the 48-hour maximum urinary excretion was approximately three times that obtained in the previous tests by the oral route, i.e. 28.0 mg for the 100-mg dose group and 37.1 mg for the 60-mg group.

c. A silica sol, prepared by diluting a commercial preparation with average particle size of 250 A° (25 μ), was given orally to two groups of five animals each: 1% (93.1 mg SiO₂) and 5% (567.0 mg SiO₂). The authors indicated that in their preliminary study the undiluted sol gelled rapidly at a pH below 9, but gelling did not occur at any pH if the sol was first diluted to a concentration of 5% or less. The urinary silica was present entirely in the "soluble" or "molybdate reactive" state indicating that the silica particles of the size present in the sol were not excreted without a prior depolymerization to the "soluble" silica form. With the group receiving the 5% silica sol containing 567.0 mg SiO₂, the net excretion was 487.2 ± 21.57 mg or ca 86% of the administered dose.

d. When 4% silica sol, prepared from the above mentioned commercial sol, was administered i.p., it again caused a large increase in urinary silica excretion over that from the oral route. When 438.8 mg were administered the first 48-hour urinary excretion period reflected the maximum of 50.8 ± 4.21 mg and returned to the normal level at the end of 20 days (10 periods). The net silica excretion for this test was 277.5 ± 20.99 mg, which represented 63% of the initial dose.

e. Precipitated silicic acid (of undetermined particle size)

was given both orally and i.p. at a level of 10 ml of suspension containing 390 mg of SiO_2 . The urinary silica levels resulting from oral administration followed a similar pattern to that of the oral silica sol described in section c above. The net silica excretion was 251.2 ± 25.80 mg, representing 64% of the initial dose. The authors postulate that the relatively low recovery may have been due to silica retention in the intestinal tract. For the i.p. injection, the net silica excretion was 187.0 ± 15.16 mg, or ca 48% of the injected dose. Again, they assume the remainder was retained by the tissues. The authors point out that after i.p. administration there was no well-defined increase in silica content of the fecal ash, and stress that the question of silica excretion into the intestinal tract after i.p. injection requires further study.

In summary, the authors interpret the above results as follows:

- (1) urinary excretion apparently was limited by restricted absorption from the gastrointestinal tract, and not by the blood transport mechanism or kidneys; in the case of the first oral tests, (section a) the silica sol prepared from the sodium silicate (section b) given i.p. in a comparable dose caused a threefold increase in silica excretion; (2) with oral administration of silica sol and precipitated silica the particles probably underwent depolymerization prior to excretion; and (3) conditions in the peritoneal cavity probably favor the depolymerization and subsequent excretion of silica. However, the reason for the greater silica urinary excretion after i.p. dosage is not known.

Finally, the authors proposed that the absorption and urinary excretion of the ingested silica is limited by its solubility in the contents of the gastrointestinal tract. They supported this view by tests which showed that when guinea pigs were dosed with 0.96 ml of tetraethyl orthosilicate dissolved in 2.0 ml of ethanol, urinary excretion values in excess of 40 mg/48 hours were obtained.

2. Settle and Sauer (445), continuing the work described in the preceding section, pointed out that when the results of the earlier experiments were considered it was noted that soluble silica in the urine did not polymerize unless the concentration exceeded 18 mg/100 ml. This indicated that the nucleus of a siliceous calculus is probably formed at some point in the urinary tract. To explore this possibility further, they treated adult guinea pigs with either 15 ml of a solution of sodium metasilicate (20 mg of SiO_2 /ml and with a pH 5 adjusted with 3N HCl just prior to dosing) i.p., or orally with 2 ml of tetraethyl orthosilicate in ethanol which contained 253.8 mg of SiO_2 /ml. The guinea pigs were sacrificed 24 hours after dosage, and kidney sections were prepared for histological study. Deposits found in the kidney tubules were confirmed as siliceous by ashing and by viewing by reflected dark-field illumination -- first, before and after treatment with 50% HCl, and then with 0.5% hydrofluoric acid. Such deposits were found in sections from both the i.p. and the orally treated groups. The authors point out that although the amount of soluble silica given was considerably greater than animals would normally ingest on a forage diet, the results do indicate that insoluble deposits may be formed in the kidneys if large enough quantities are given.

C. Rabbits

It is appropriate at this point to mention the work of Collet and Moussard in regard to biliary elimination of silica (103). By anesthetizing the rabbit and catheterizing the common bile duct, the bile was collected and biliary output recorded on a rheograph. Control bile was collected for 30 minutes and then the animal was injected with 1.25 cc/kg of a suspension of highly dispersed silica in distilled water into the saphenous vein. (The sample was assumed to contain a mixture of silicic acid monomers and polymers). The silica content of the samples was determined by a modified version of the method of King (check original version of 103). The biliary elimination of silica in relation to the quantity injected i.v. was considered negligible by the authors, i.e. ranging from 0.35 to 1.1% during the 6 hours following injection.

D. Dogs

1. Particulate and soluble silicas, as well as silicic acid and organic silicate (Silistren), have been administered to dogs (260) and the excretion followed. Pulverized quartz was given to dogs by stomach tube, as shown in Table 21, with the excretion pattern as indicated there. Analyses were accomplished by the procedure of King and Stantial [Biochem. J. 27: 990-1001 (1933)].

Table 21 (260)

Excretion of Silica in the Urine after Administration by
Stomach Tube of 5 g Powdered Quartz, Suspended in Water

Hour	SiO ₂ per 100 cc. urine	SiO ₂ per 100 cc. blood
	mg.	mg.
0	2.7	0.8
4	14.3	--
7	14.7	0.7
16	4.8	--
24	3.3	0.7

Soluble silicas -- for example solutions of sodium silicate -- could not be administered in their normal alkaline solution because dogs would immediately vomit the material. However, when they were neutralized to phenolphthalein with dilute hydrochloric acid just prior to dosage by stomach tube, the dogs tolerated the dosage well. In the stomach, deposits of gelatinous silicic acid no doubt tended to form from the 5% sodium silicate solutions, when neutralized thusly, and these were then somewhat redissolved by the alkaline juices of the small intestine. The authors believe, however, that a large part of the silicate administered this way was precipitated as silicic acid before it had been in the stomach long, and, although a portion was redissolved, a large amount still remained out of solution but was absorbed and eliminated quickly in the urine. Table 22 illustrates the results of various dose levels of sodium silicate administered as described.

Silica Content of the Urine after Administration of Silicic Acid by Stomach Tube (1) and Ingestion (2)

The authors state that analyses of the organs of several of the dogs showed no increase in silica content over controls -- a finding that contradicts those of other workers (433,511). However, as already discussed, the analytical methodology may have been a limiting factor.

74

elevation. The authors gave data showing that only 50% of the administered dose was recovered in the urine, and they postulated that in the case of these i.v. dosages some was probably excreted in the feces with a fairly large proportion probably retained in the body. Spleen analysis of one dog confirmed a considerably higher than normal silica content.

"Silistren", a glycol ester of silicic acid, was administered to dogs as an example of organically bound silica. An amount of silistren in water solution corresponding to 400 mg of silica was injected i.v. in two dogs. In one dog, urinary silica reached almost as high a level as that obtained with silicic acid when given by stomach tube, but in the other a more moderate increase was obtained; bloody urine and anuria resulted in the latter case. Blood silica was raised and maintained at a higher level with Silistren than with any other means tried. The authors suggest that, when present as an organic compound, the silicic acid can remain at high concentrations when it is incapable of remaining there as inorganic silicate.

E. Humans

1. Page, Heffner and Frey (380) measured the urinary excretion of silica in healthy young males following the administration of magnesium trisilicate (Tabliod - a commercial antacid tablet, which readily disintegrates in water). Five grams were given daily in five 1 gram spaced doses for four consecutive days. Twenty-four-hour urine excretion was collected for 5 days and each specimen analyzed for silica. Table 23 reports the results of these analyses.

Table 23 (380)

Experimental Study on Same Subjects as Table 21 with Daily
Oral Ingestion of 5 grams of Magnesium Trisilicate
(Urinary Silica (SiO_2) -- 24 Hour Totals and per 100 cc. Urine)
--SUBJECTS--

Day	Mg. Tri. Orally	S		R		M		L		B		Mean Tse.
		Total SiO_2 (Mg.)	Mg./100 cc.	Total SiO_2 (Mg.)	Mg./100 cc.	Total SiO_2 (Mg.)	Mg./100 cc.	Total SiO_2 (Mg.)	Mg./100 cc.	Total SiO_2 (Mg.)	Mg./100 cc.	
1	5 Gm.
2	5 Gm.	152.0	14.6	167.0	9.09	191.0	13.1	196.0	20.0	151.5	8.51	172.1
3	5 Gm.	200.0	20.0	118.0	8.60	202.0	9.00	230.0	19.0	141.0	9.01	178.5
4	5 Gm.	203.0	10.4	152.0	10.0	152.0	10.0	114.0	8.30	162.7
5	None	96.0	10.5	40.6	2.78	89.0	9.00	45.6	3.8	50.0	3.6	64.0
6	None	33.0	2.4	25.8	2.0	11.8	1.0	24.5
Total excreted from 2nd to 6th day inclusive.		448.0		528.6		637.0		653.6		459.5		545.3
Expected excretion (calculated from mean normal daily output x number of days).		*47.4		72.8		76.0		56.8		55.2		61.6
Difference =		X 400.6		455.8		561.0		596.8		404.3		483.7

Mg. Tri. = Magnesium Trisilicate. Tse. = Total SiO_2 Excreted.

As can be noted from the table each subject took a total amount of 20 grams of magnesium trisilicate, which contained ca 9.2 grams SiO_2 . The increased SiO_2 excretion from the second day until three days after the end of treatment averaged 484 mg, so that ca 5.2% of the ingested silica was excreted in the urine. No deleterious effect was noted from the ingestion of the magnesium trisilicate.

2. A case of urinary calculi containing silica (302), as described by Lipworth, Bloomberg, and Reid, is one of the very few reported in the literature. A 62-year-old man, with hiatus hernia

and resultant pyrosis had taken magnesium trisilicate for years, 2 to 4 times daily. He also had coronary artery disease, with two or probably three attacks of coronary thrombosis, and had had anti-coagulant therapy; he was hypercholesteremic and for some years had been on a diet low in dairy and animal fats and high in fruit and vegetables. In June 1961 the man complained of pain and a calculus was diagnosed in his left lower ureter with blockage. He passed some calculous fragments within the next few days. Again in October of that year he passed "gravel" which was found to consist almost entirely of inorganic matter. Tests for urates, oxalates and calcium were negative, but there was insufficient material for further analysis. In June of the next year, after prostatectomy, he passed a calculus presumed from his description of pain also to have come from the left ureter. It was irregularly round, facettted, ca 0.5-1 cm in diameter, and brown in color. Calcium, magnesium, oxalate, uric acid, cystine, and xanthine were not detected. A trace of phosphate and slight amount of organic matter were present, and, upon further analysis, the material was found to contain ca 50% silica. The author points out that although the pathogenesis is obscure, and only eight previous cases had been reported in the literature, there is a possibility that silica calculi may not be a rare complication.

3. Five of the eight cases of urinary calculi mentioned by Lipworth et al. (section above) were reported by Lagergren (288). The latter author discussed the survey of some 800 patients who had passed urinary calculi. The stones were analyzed by X-ray crystallography and it was found that five of the patients passed calculi

which consisted mainly of silica, present in a finely crystalline form. All five patients had been taking magnesium trisilicate in tablet form regularly, over a period of several years, as an antacid for gastritis or ulcer. There was no evidence that any of the patients had exceeded the manufacturers' specified doses.

4. Langendorff and Lang (291) studied the effect of polymeric silicic acid on renal SiO_2 excretion in man. Aerosil^R, as described by these authors is an amorphous aerosol with particle size between 10 and 40 μm and a surface of $175 \text{ m}^2/\text{g}$, having a surface free of pores and with siloxan groups and silanol groups ($5\text{--}6 \mu\text{mol OH-groups}/\text{m}^2$) thus capable of forming hydrogen bonds. One g of aerosil is capable of taking up ca 400 mg water by multi-layer adsorption without losing the quality of a dry powder. Ca 9 mg is fixed to the surface by hydrogen bonding. Also tested were a beer-clarifying agent FK 700, which is a pure hydrated silicic acid containing 86.65% SiO_2 . Ten men and two women, between 22 and 28 years of age, were given Aerosil^R or FK 700, two doses of 1.25 g in 250 ml of commercial applejuice (total dose 2.5 g), on the 4th day of a 7-day period. Twenty-four-hour urine specimens were collected on days one to seven and samples from each were frozen until analyzed for SiO_2 content. The determinations were made by the method of Bauman [Z. physiol. Chemie 319: 38 (1960)], with an attempt to determine also polymeric SiO_2 (SiO_2 determination after alkaline hydrolysis). As a rule, somewhat higher values were found for "total SiO_2 " but these fell within the limits of error of the method. This fact tends to support the finding of Settle and Sauer (445) that soluble silica in the urine

does not polymerize below 18 mg/100 ml. Table 24 gives the results of these analyses.

Table 24 (291)

Aerosil® Preparations

Tag	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
Vpn.-Nr.	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
1	0,80	26,2	33	1,78	85,1	48	0,93	64,7	67	2,01	51,8	26	1,05	14,0	13	1,39	34,8	25
2	0,77	28,1	36	1,84	119,8	65	1,07	92,0	86	1,62	23,6	15	0,76	24,6	32	0,75	32,5	43
3	0,98	49,3	50	1,20	55,7	46	1,28	31,7	25	1,13	45,9	41	0,75	37,3	50	1,08	41,2	38
4	0,73	30,0	41	1,91	75,2	39	0,81	61,9	76	1,69	83,9	49	0,89	35,7	40	1,05	39,9	38
5	0,69	50,0	72	1,50	56,0	37	2,40	101,5	42	1,65	66,1	40	1,21	49,1	41	1,63	65,3	40
6	0,63	23,6	37	1,15	47,7	41	0,97	34,6	36	0,82	35,1	43	0,67	57,5	80	0,83	41,5	50
7	0,54	25,7	48	2,27	78,9	35	0,72	47,0	65	1,57	58,1	37	1,00	35,5	30	1,14	64,8	57
1.-3.	2,55	103,6		4,82	200,6		3,28	188,4		4,76	121,1		2,56	75,9		3,22	108,5	
4.-7.	2,59	129,3		6,83	257,8		4,90	245,0		5,73	243,2		3,77	177,8		4,65	211,5	
Mittelwert 1.-3.	0,85	34,5	41	1,61	86,9	54	1,09	62,8	57	1,59	40,4	25	0,85	25,3	29	1,07	36,2	34
Mittelwert 4.-7.	0,65	32,3	49	1,71	64,5	37	1,23	61,3	50	1,43	60,8	42	0,94	44,5	47	1,16	52,9	45

4 Spalte 1: Harnvolumen, Liter; Spalte 2: tägliche SiO₂-Ausscheidung in mg.
Spalte 3: SiO₂-Konzentration in µg/ml.

FK 700 Preparation

Tag	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
Vpn.-Nr.	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
1	0,94	46,4	49	0,95	34,3	36	1,01	66,6	66	1,43	17,6	12	1,07	33,0	31	0,75	33,7	45
2	0,76	50,8	67	0,72	33,9	47	0,94	73,1	78	1,28	16,2	13	1,32	49,1	37	0,85	27,4	32
3	0,74	30,0	40	1,33	42,2	32	1,10	74,6	68	1,46	14,5	10	1,63	37,7	23	0,55	21,0	38
4	0,85	53,3	63	1,81	55,8	31	1,20	48,9	41	1,77	24,6	14	1,45	44,7	31	0,96	28,8	30
5	0,90	62,7	70	1,77	61,8	35	2,58	142,3	55	0,87	19,5	22	0,81	45,6	56	0,68	26,6	39
6	0,72	50,7	70	0,90	75,2	84	0,83	75,7	91	1,20	20,6	17	0,93	44,0	47	0,60	11,7	19
7	0,98	42,0	43	1,08	35,1	33	0,86	58,2	68	1,46	17,0	12	1,55	98,8	64	0,53	19,9	38
1.-3.	2,44	127,2		3,00	110,4		3,05	214,3		4,17	48,3		4,02	119,8		2,15	82,1	
4.-7.	3,45	208,7		5,56	227,9		5,47	325,1		5,30	81,7		4,74	233,1		2,77	87,0	
Mittelwert 1.-3.	0,81	42,4	52	1,00	36,8	37	1,02	71,4	70	1,39	16,1	11	1,34	39,9	30	0,72	27,4	38
Mittelwert 4.-7.	0,86	52,2	60	1,39	57,0	41	1,37	81,3	59	1,33	20,4	15	1,19	58,3	49	0,69	21,8	31

4 Spalte 1: Harnvolumen, Liter; Spalte 2: tägliche SiO₂-Ausscheidung in mg.
Spalte 3: SiO₂-Konzentration in µg/ml.

1. Day
2. Experimental subject number
3. Mean value
4. Column 1: urine volume, liter;
Column 2: daily SiO₂ excretion in mg;
Column 3: SiO₂ concentration in µg/ml

As can be expected, and also noted from the table, excretion varies greatly between individuals and from one day to the next in the same individual, possibly due to variations in diet or liquid intake. For the group, during the preliminary or control period, the SiO_2 concentrations were between 11 and 70 $\mu\text{g}/\text{ml}$ with average daily excretion between 15.1 and 86.9 mg SiO_2 . After the administration of 2.5 g Aerosil^R the SiO_2 excretion did not clearly change. In some individuals there was a reduction after one day of treatment, while the others showed an increase in SiO_2 excretion. With FK 700 (2.15 g SiO_2) there was a somewhat clearer picture. SiO_2 excretion increased in all subjects on the fourth day of treatment, with the exception of one whose SiO_2 excretion decreased 26 mg. However, the author points out that in general, in both groups, neither the increased excretion in individual cases nor the total SiO_2 excretion during the four days following the administration was related to the quantities of 2.5 or 2.15 g SiO_2 administered. No mention was made as to the possible fate of the bulk of the SiO_2 -- absorption in the tissues or excretion in feces.

Various values have been estimated for the maximum limit of silicic acid in the monomeric state, when in water or urine: Settle and Sauer (445), 18 mg/100 ml; Thomas [Dtsch. Z. Verdauungs- u Stoffwechselkrankheiten 25: 260 (1965)], 120 $\mu\text{g}/\text{ml}$ (12 mg/100 ml); and Bauman [(Z. physiol. Chemie 320: 11 (1960)], 600 $\mu\text{g}/\text{ml}$ (60 mg/100 ml). However, the SiO_2 concentrations found in urine in these experiments are below the range in which polymeric SiO_2 harmful to the urinary tract is formed.

5. In an effort to determine if a relationship exists between urinary silica excretion and silicosis, Whitehouse (521) administered four different types of dust as well as silicic acid, to humans by mouth. Administered in suspension with 200 cc of water about half an hour before the mid-day meal were: quartz, with particle size $6\ \mu$; flint of the same size; scotch whinstone of the same size (described as a rock containing no free silica, silicates being present chiefly as labradorite, augite and serpentine); and precipitated silica obtained from sodium metasilicate in the usual way. A dose of 0.2 g of silicic acid (the soluble silica) was also taken by mouth. The urinary output of SiO_2 from these substances is shown in Table 25.

Table 25 (521)

The Effect of Oral Administration of Silica on
the Excretion of Silica in Urine

REPORT- MENT NUMBER	DOSE TAKEN BY MOUTH	DATE	VOLUME OF URINE	SPECIFIC GRAVITY OF URINE	SiO_2 IN URINE	TOTAL SiO_2 IN 24 HRS. URINE
			cc.		mgm./100 cc.	mgm.
I	2 gm. whinstone	16/ 5/35	1724	1.016	0.85	14.7
		17/ 5/35	1718	1.015	0.73	12.5
		22/ 5/35	1506	1.019	0.90	13.6
		23/ 5/35	1297	1.019	1.27	16.5
II	1 gm. quartz	29/ 5/35	1545	1.018	0.78	12.1
		30/ 5/35	1387	1.019	0.62	8.6
III	3 gm. quartz	5/ 6/35	1783	1.014	0.61	10.9
		6/ 6/35	1610	1.017	0.67	10.8
		7/ 6/35	1626	1.016	0.43	7.0
IV	6 gm. quartz	17/ 6/35	925	1.023	1.28	11.8
		18/ 6/35	1596	1.016	0.74	11.8
V	0.2 gm. silica as silicic acid	27/ 6/35	1850	1.014	0.70	13.0
		28/ 6/35	1590	1.020	7.82	124.4
VI	9.3 gm. quartz	3/ 7/35	1345	1.018	0.78	10.5
		4/ 7/35	1595	1.014	1.12	17.8
		5/ 7/35	1270	1.019	0.83	10.5
VII	0.5 gm. precipitated silica	1/ 8/35	1480	1.020	0.82	12.1
		2/ 8/35	2117	1.019	0.67	14.2
VIII	2.5 gm. precipitated silica	22/ 8/35	1095	1.026	1.44	15.8
		23/ 8/35	1485	1.020	2.40	35.7
		30/ 9/35	1775	1.018	0.63	11.2
IX	6 gm. flint	1/10/35	1985	1.019	0.55	11.0
X	6 gm. whinstone	28/10/35	1650	1.015	0.88	14.5
		29/10/35	1855	1.014	2.40	44.5
XI	0.22 gm. flint*	1/ 4/36	1260	1.017	0.68	8.6
		2/ 4/36	1545	1.016	1.07	16.5
XII	2.28 gm. flint*	8/ 9/36	1720	1.013	0.76	13.1
		9/ 9/36	2040	1.014	1.38	28.1

* Maximum particle size $1\ \mu$.

It was recognized that for positive or increased urinary SiO_2 excretion either the dosage must be increased or the range of particle size decreased, and this was undertaken. Flint dust of 1 μ was prepared by water sedimentation and administered in water as in the first series of experiments. Results from this study, along with additional data on separate samples of urine from individuals receiving precipitated silica, silicic acid, and whinstone are shown in Table 26.

Table 26 (521)
Urinary Excretion of Administered Silica

EXPERI- MENT NUMBER	DOSE TAKEN BY MOUTH	SAMPLE	PERIOD OF URINE COLLECTION	VOLUME OF URINE	SPECIFIC GRAVITY OF URINE	SiO_2 IN URINE	TOTAL SiO_2
			<i>hrs. after dose</i>	<i>cc.</i>		<i>mgm./100 cc.</i>	<i>mgm.</i>
V	0.2 gm. silica (as silicic acid)	(a)	0-5	452	1.015	17.52	70.2
		(b)	5-12	560	1.018	5.72	32.0
		(c)	12-21	490	1.023	2.28	11.2
		(d)	21-24	88	1.024	2.28	2.0
VIII	2.5 gm. precipitated silica	(a)	0-4	246	1.030	6.03	14.8
		(b)	4-10½	313	1.028	4.74	14.8
		(c)	10½-12	114	1.023	1.68	1.9
		(d)	12-22	753	1.017	0.45	3.4
		(e)	22-24	59	1.026	1.43	0.8
X	6 gm. whinstone	(a)	0-4	350	1.020	5.42	19.0
		(b)	4-8½	1080	1.012	0.85	9.2
		(c)	8½-12	227	1.023	4.02	9.1
		(d)	12-24	198	1.029	3.63	7.2
XII	2.28 gm. flint	(a)	0-1½	680	1.011	1.04	7.1
		(b)	1½-4	250	1.015	2.77	6.9
		(c)	4-8	400	1.014	1.12	4.5
		(d)	8-12	225	1.028	2.39	5.4
		(e)	12-24	485	1.018	0.86	4.2

With the larger dose of the fine-particle-size flint, a definite increase in urinary silica excretion was noted; the same was true for precipitated silica and, to a greater extent, silicic acid. In the case of the Scotch whinstone the daily output of urinary silica was increased to at least three times the normal figure -- a result the author did not expect in view of the lack of free silica in the rock. However, on the basis of solubility studies on the substances tested, he concluded that solution of the whinstone had been caused by the action of the acid gastric juice. The author reminded readers that variations in the silica content of different water supplies should be taken into account in considering relative values, and also diet, particularly if silica is ingested in such a state that it is readily absorbed -- e.g. beer, water etc.

IV. Effects on Enzymes and Other Biochemical Parameters

A somewhat general condition, while not of extreme or necessarily permanent nature, has been shown to be adversely affected by the administration of a drug containing silica. Rohm, Seybold and Pirtkien (420) have discussed the induction of ulcers in the antestomach in rats and the influence of ten different drugs commonly used in human gastric therapy. Male Wistar albino rats were treated with a 20% glucose solution administered from drinking tubes; the treatment lasted for eight days. The various drugs were in part dissolved in the glucose, and in part injected s.c., i.m. or i.p. The rats were sacrificed and the stomach carefully removed for histological examination, with the ulcers being evaluated according

to number and size. Of the ten drugs, only one appeared to inhibit ulcer formation, and this conclusion was not statistically confirmed. Five, however -- among them aluminum sodium silicate with 0.5% belladonna extract (Belladonna-Neutralon) -- produced a statistically confirmed increase in ulceration.

A. Renal Mechanisms and Lesions

1. In an often quoted work, Policard and Collet (398) described their experiments with the injection of strong concentrations of fresh silica gel into the peritoneal cavity of albino rats. The silica gel was prepared by electrolysis of sodium silicate, which yields a very pure product with silica concentrations ca 5%. This is a polymer of silicic acid $\text{SiO}_2 \cdot n\text{H}_2\text{O}$ with a consistency of a semi-solid gel. It was sterilized with UV rays and chemically analyzed to establish a volume to weight ratio for SiO_2 . The gel was administered, i.p. 45 to 90 mg of silica, in a single dose. The animals were sacrificed or died spontaneously in from 16 hours to 20 days. Kidney sections were prepared for histological study and the following observations then made. Sixteen hours after injection a typical acute toxic epithelial nephritis was noted, recalling mercury nephritis, without glomerular involvement. At about the seventh day, a transitional period was noted during which the toxic nephritis subsided and foci of interstitial nephritis appeared. As a rule this period was marked by a large number of inflammatory lesions of the pyelonephritic type, with highly dilated tubules. After this period came a phase of progressive tubular atrophy and interstitial fibrosis and glomerular alterations which the authors term interstitial

glomerulonephritis.

2. In a later study from the same laboratory the effects of oligomeric silicic acid were considered (399). This acid was found capable of penetrating by diffusion into the mitochondria of the renal epithelium of rats that had received i.p. silica gel some hours previously. Inside the mitochondria, where a different equilibrium medium existed, the silica polymerized and became visible in the high resolution microscope in the form of very fine, very dense granulations measuring some 40 \AA (4μ). These granulations were joined together and formed very dense heterogeneous granular masses of a size varying between ca 200 and 500 \AA ($20\text{--}50 \mu$). They were always situated between the widely spaced internal membranes of the mitochondria. After the development of these silica deposits, the mitochondria do not remain normal. Two types of conditions progress. First, the mitochondria undergo swelling, with internal membranes becoming widely separated and progressively fainter. The silica masses increase and gradually the outlines of the mitochondria blur. They appear to break up in the surrounding cytoplasmic milieu which is much altered by edema; the masses of silica are then free in the protein magma resulting from the degeneration of the renal epithelial cells. This magma occupies the lumen of the tubules and constitutes the well-known "granular casts." This swelling does not seem to be directly dependent upon the presence of the silica masses however, as the adjacent mitochondria without silica masses may reveal identical alterations. The swelling appears to be related to a condition of the ambient cytoplasmic milieu, which could possibly be the presence

of diffusible oligomeric silica in the cytoplasm. In a second type, instead of being swollen the mitochondria become denser without changing dimensions. The light and swollen mitochondria and the denser mitochondria are not mixed together, but seem to occur in adjacent zones of the degenerated epithelium. The two types of mitochondria, together with their silica masses, reappear in the casts that occupy the lumen of the injured urinary tubules. These casts then are carried by the glomerular urine and are excreted. In the urinary tubules cleared in this way, the regeneration of a new epithelium may take place. The authors compare this circuit of elimination with the purification of the lungs where the degenerated alveolar cells, progressively eliminated through the bronchial passages, rid the lungs of silica particles. In neither case, kidney or lung, is the purification process entirely complete, and this is the cause of persistent and often progressive sclerotic foci (pneumoconiosis or sclerous silicotic nephritis).

3. Markovic and Arambasic (329) induced chronic interstitial nephritis in guinea pigs by administering pure silica (quartz), of particle size 1-3 μ in diameter, in drinking water for a period of several months. The animals developed renal pathological changes similar to those found in man in Yugoslavia. Endemic nephropathy was attributed to rock erosion in village communities on the banks of the lower reaches of large rivers. According to the authors, the renal lesions observed in such a situation were caused by release of silicic acid, and specifically involved a dystrophic-atrophic lesion in the parenchyma and a proliferative-inflammatory lesion

in the interstitial tissue.) The lowest concentration of silicate suspension that induced interstitial nephritis in the animals was 50 mg/liter of water -- a quantity of SiO_2 detectable in the drinking water in areas where human endemic nephropathy was observed. Also, the authors point out that the human kidney is more sensitive to silicate than the kidney of herbivorous animals because of the acid pH of the human urine.

B. Serum Protein and Protein Biosynthesis Alterations

1. Amorphous and crystalline silicon dioxide have been studied for their effect on blood protein fractions and histamine (186). Gel'fon and Fedorova treated white rats, M, BW 250 to 300 g with 50 mg of silica gel dust suspended in 1 ml of physiologic saline, injected intratracheally. A second group was treated similarly with 50 mg/ml of crystalline silicon dioxide dust also in physiologic saline. Particle sizes were 1 to 3 μ . The animals were sacrificed 1, 2, 4, and 7 months after injection. Serum proteins were determined by paper electrophoresis. The administration of crystalline silicon dioxide resulted in a slightly elevated total protein level in the serum, decrease in albumin, and increase in the globulins (mainly β - and γ - globulin fractions). In this group histamine increased in the lung tissues. In the amorphous-silicon-dioxide treated groups, the trends were the same but less pronounced, and abnormal values occurred later than with the crystalline group. The lungs contained cellular dust foci unlike the silicotic nodules typical of experimental silicosis in white rats, focal thickening of the septa, signs of emphysema, and slight peribronchial and perivascular

sclerosis. In rabbits injected s.c. with quartz dust, the serum albumin level was lowered, and there was a gradual increase in the amount of globulins (γ -globulins in particular) and in blood histamine. Quartz dust, when administered to rats who had been previously immunized with typhoid vaccine caused greater changes in the serum albumin content and the β - and γ -globulins than when administered to nonimmunized animals. However, the morphological changes in the lungs and lymph nodes were almost identical in both groups of animals.

2. An in vitro study of silica effect on protein biosynthesis by rat-liver cell-free systems (105) has shown that leucine incorporation into microsomal protein was depressed by both crystalline HF-treated silica (tridymite) and vitreous (HF-treated) silica. As can be seen from Table 27, the effect was more pronounced with the crystalline form, a circumstance which the authors interpret as probably related to the surface structure of the silica particle.

Table 27 (105)

In Vitro Incorporation of DL-leucine-1-C¹⁴
into Rat Liver Microsomal Protein.

Each figure (c.p.m. per mg protein) is the mean \pm S.E. of 2 flasks

Experiment	Control	Crystalline silica	Vitreous silica
1	92.5 \pm 4.47	52.0 \pm 1.25	..
2	80.5 \pm 5.48	59.0 \pm 5.00	..
3	116.0 \pm 3.00	62.0 \pm 1.40	..
4	111.0 \pm 9.00	65.0 \pm 6.00	..
1	81.0 \pm 3.00	..	65.5 \pm 7.18
2	75.5 \pm 10.19	..	13.0 \pm 6.00
3	84.0 \pm 10.00	..	76.5 \pm 3.46
4	80.5 \pm 5.18	..	62.5 \pm 7.18

The statistical evaluation of data has been made with the analysis of variance test.

3. Repeated intravenous injections of 1-5 μ silica in 100 mg doses were given to rabbits (359), with the total dose of 500-600 mg spread over a 10-15 month treatment period. The proteinograms, developed by electrophoresis, revealed essentially an imbalance in the albumin-globulin ratio in favor of the globulins. There was a distinct increase in all cases in the β - and γ -globulins, a phenomenon which agrees with the observations of Gel'fon (186) on rats, Mosinger (357) on rabbits and Autio (045) on dogs. In this study (359), there was also an increase in the α -2-glycoglobulins and β -lipoproteins. Furthermore, there was a reduction in urinary 17-ketosteroids, ca 50%, in both male and female animals. Histologically, in the liver a non-specific cirrhotic process was observed with nodular silicosis. Silicotic lesions were also seen in the spleen, lymph nodes, and lungs. The pulmonary silicotic nodules invariably had a septal topography and always developed around pulmonary capillaries. They were accompanied by pulmonary emphysema. In the kidneys there was always glomerulitis and tubular nephritis. Finally, diffuse amyloidosis was observed, with diffuse calcinosis; aortic lesions; and general mesenchymal, visceral and neuro-endocrine reactive syndrome.

C. Blood Lipids

Szabo, Mody and Szekely (481) have reported on the effect of silicic acid and silica dioxide on blood lipids. Twelve rabbits were injected s.c. 5 times in a period of 11 days with 0.5 ml/kg of an 0.5% solution of colloidal silicic acid. They also received 20 mg cortisone i.m. daily. Serum proteins and lipoproteins were determined

before (for control base) and one and two weeks after the first injection. Every animal developed severe dysproteinemia. The blood lipid level rose and the relative and absolute amounts of the β -lipoproteins increased, as did the β/α ratio. Cortisone had no effect on the changes. Heparin (400 units) restored the normal blood-lipid and β -lipoprotein levels. In another series of tests, 14 dogs were injected i.v. every 2 or 3 days over a period of 30 days with 2 ml/kg of the same colloidal silicic acid solution. Again serum proteins and lipoproteins were determined before and every 10 days after injection. The amount of lipoproteins was high in all cases and the β/α ratio markedly increased. Guinea pigs, in the third test by these authors, were studied for the effect of silicon on serum cholesterol level. Twenty-six pigs were given a single i.p. injection of 50 mg of silicon dust or 10 mg of amorphous silicon dioxide. The blood cholesterol level rose significantly on the average to 118.1 ± 8.7 and 109.6 ± 9.3 mg% respectively, the control value being 84.7 ± 2.4 mg%.

D. Enzyme Changes

1. Talc has been observed to cause a decrease in hepatic catalase activity (245). Strong A mice were given s.c. injections of 1.0 ml of a 0.67% suspension of talc. A transient, but marked, rise in catalase activity was observed on the 4th day following injection. The rise in activity was manifest mainly in the mitochondrial fraction, with concomitant increased incorporation of radioactive leucine in the microsomal fraction. The authors propose a relationship between the increased catalase activity and increased incorporation of

leucine. Attention is also called to the in vitro study of silica effect on protein biosynthesis by the rat liver (105) discussed on page 88 of this monograph.

2. Webster et al. (513) have studied some of the biologically active substances produced in the state of "silica shock" in rabbits and guinea pigs. Using Aerosil [of particle size 100-400 Å (10-40 mμ)] and ground silica of particle size <1 μ, suspensions were made to contain 25 mg in 2 ml which were injected into a vein in one ear. Analysis of serum of the animals in the resulting state of shock indicated an increase of a kinin, probably bradykinin, and the authors suggest that the release of a kinin may be a factor in the cause of death in a state of silica shock.

E. Effects on Blood Cells, and Clotting

1. Zaks and Galeniyetse (537) have measured the heparin number of blood and heparinocytes after oral administration of silica gel to 11 healthy humans. They received silica gel, 2.5 ml of a 20% sodium silicate solution neutralized with normal HCl acid solution, for 21 days. After receiving the gel, the humans exhibited an increase in the heparin number, the first symptoms appearing about the 7th day. A statistically significant increase in the heparin number was observed on the 21st day. The number of basophilic leukocytes decreased on days 7 and 11 and increased on days 14 and 21, but did not reach the baseline level.

2. The effect of colloidal silica on blood clotting was reported by Margolis (327). He found that the silica accelerates blood coagulation by adsorption and partial denaturation of a specific

plasma protein, the Hageman factor. Quantitative measurements showed that the activity of the silica depends on the particle diameter, and indicates that somewhere between 10 and 2 μ there is a critical minimum particle size below which silica ceases to activate plasma.

3. The hemolytic effects of colloidal silica (217) and aerosil, crystalline silica, silicic acid powder, and silica gel (316) have been studied. A detailed discussion of the in vitro studies on human cells (217) and sheep erythrocytes seems beyond the scope of this monograph. However, it is appropriate to point out that Margolis (217) indicates that this phenomenon, as with the effect of colloidal silica on blood clotting, may be related to the geometrical relation between the colloidal particles and the protein molecules.

V. Drug Interaction

1. The now well established phenomenon of intraperitoneal adhesions due to talc has been considered in relation to cortisone effect (056), with what the authors consider to be definitive results. A total of 89 talc-injected mice were each given s.c. 0.5 mg cortisone daily from the day of talc injection until sacrifice. Talc was administered 35 mg i.p., as 0.35 ml of a 10% suspension. In a second series, 13 additional mice were given cortisone daily only during the first 10 days, and 13 more daily only during the last 10 days. In a third series eight mice were pretreated daily with cortisone for 14 days before talc injection, and then daily for an additional seven days. Histological examination showed that although cortisone delayed the onset of formation of abdominal adhesions due to i.p.

injection of talc after the adhesions had been formed, cortisone had no effect in promoting their resolution.

2. Also, the influence of cortisone on leptomeningeal reactions induced by talc has been studied (379). Eighteen rabbits received intrathecal injections of 1.0 cc of talc suspension with sacrifice in 15 days. Ten of these animals were treated with daily injections of cortisone 5 and 10 mg/kg. The control (talc only) animals developed a meningeal reaction consisting of the appearance of histocytes, macrophages and giant cells, while the cortisone treated rabbits showed a marked diminution of the meningeal response to the talc. Preliminary experiments with the use of hydrocortisone, injected intrathecally (10 mg) on the 1st and 7th days also suggest diminution of the meningeal response to talc, though not as strikingly so as with i.m. cortisone.

3. Silicon compounds have been noted to have an effect on the formation of atheroma (platelet deposits of cholesterol), and Loeper and Loeper (305) have investigated the use of sodium silicate, lysine silicate, and the organic silicon compounds, silanol salicylate and silanol citrate. The control rabbits were kept on an atherogenic diet that included 0.75 g/day of cholesterol dissolved in oil, while other groups were kept on the same diet while receiving:

(a) sodium silicate i.v. in a dose of 0.01 g on alternate days, (b) lysine silicate p.o. 0.01 g/day, and (c) silanol salicylate (monomethyltrisilanol salicylate) i.v. 0.0075 g on alternate days or (d) silanol citrate (monoethyltrisilanol citrate) i.v. 0.02 g on alternate days. After sacrifice at two months, gross examination

of the entire aorta was made and histological examination with lipid staining was carried out. The results are shown in Table 28.

Table 28 (305)

Effect of Silicon Compounds on Experimental Atheroma

Diet	Rabbits	Atheromatous rabbits	
Cholesterol	8	8	
Cholesterol + sodium silicate	8	3	
Cholesterol + lysine silicate	5	2	
Cholesterol	8	5	
Cholesterol + silanol salicylate	5	0	
Cholesterol + silanol citrate	4	1	

With regard to the histological examination, the authors noted that in addition to atheromatous plaques in the controls, the fine fragmented elastic fibers were frequently altered (in 15 of 22 rabbits), and there was increased metachromasia of the basic substance in the subintimal region and the media. In the treated animals, however, alteration of the elastic fibers was rare (6 of 23 animals) and, interestingly, the elastic fibers not only were intact but often were more numerous and thicker in the three aortic layers and were without collagen proliferation. This reaction was particularly marked in organic silicon compounds. Metachromasia of the basic substance was increased in both treated animals and controls. The

authors discussed the theory that the effect of silica compounds might be due to increased impermeability to lipids, but concluded that it seemed more likely it is linked to the protection and development of the elastic fibrils of the aortic wall.

4. Several other examples of the interaction of silica compounds with drugs can be found in the works of Grelet (206), Daniel-Moussard and Collet (129), and Marks (331). Grelet (206) investigated the role of vitamin A deficiency in intraperitoneal sclerosis induced by kieselguhr in rats, and found that there was no effect on the normal stages of histiogenesis of the intraperitoneal granulomas formed upon contact with the silica-containing substance. Daniel-Moussard (129) investigated the role of certain chelating agents, hoping that they would be found to "detoxify" the harmful silica-containing mineral particle. Disodium versenate, monocalcium versenate (ethylene-diamine tetracetate), B.A.L., (British Anti-Lewisite or 2-3-dimercaptopropanol), and Apresoline from CIBA, (1-hydrazinophthalazine chlorhydrate) were studied for their effect on rats treated with quartz of particle size 3 μ . In doses that were non-toxic to the animals none of the chelates had any effect on inhibiting silica fibrosis. In conclusion, Marks (331) studied 11 drugs, nine of them considered as histamine-releasing agents, while two were phenazines. The drugs were found to exert a protective effect on phagocytic cells in tissue culture against injury by silica dust, but the authors interpreted the effect as not associated with the histamine-releasing activity of the compounds. The phenazines also neutralized the toxicity of silica dust.

VI. Mechanism of Toxicity and the Factors Affecting It

The detailed mechanisms of silica toxicity were not fully described in any of the reports found. Silica can be absorbed by ingestion or by inhalation. When ingested, silica is acted upon by the stomach and intestines to form silicic acids and soluble silicate (380). Some silica is absorbed by the digestive tract and enters into the blood stream. One author states (511) that silica is generally assumed to circulate in the blood in the colloidal form. Accumulation occurs in varying degrees in some of the organs (084, 511, 273 and others) with excretion by the kidneys. A numerical estimate, made by Webb et al. (511) (on rats fed chronic doses), indicated excretion by the feces in the range of 95% of the dose administered, 4% in the urine, with 1% being accumulated in the tissues. When silica enters the body through the lungs a considerable amount is caught in the capillary bed of the lungs; while some progressively disappears with storage in other organs, a sizeable amount still remains (084 and others). The capillary bed of the lungs can also act as a filter for silica particles which have entered the blood

stream by other routes (359). Webb (511) states he does not find it surprising that the silica is picked up predominantly by organs containing a large amount of reticulo-endothelium tissue — i.e. liver and spleen.

Concerning acute toxicity, it was shown very early by Gye and Purdy (209) that massive intravascular clotting occurs in rabbits receiving i.v. injections of silicic acid (209). The authors pointed out that although this clotting was secondary to endothelial damage, it was the immediate cause of death. It was shown by Modell and Salzman (348), however, that even with administration of an anticoagulant, death occurred in cats given similar doses of silicic acid i.v. (See Table 4). Also the latter authors showed that there was no correlation between toxicity and either the acidity of the injected solution or the rate of injection. Others suggest that silicic acid exerts its toxic effect by constricting the blood vessels of the lung; and Filley, Hawley, and Wright (161) have described the specific bronchoconstriction produced by colloidal silica.

Much emphasis in the literature has been placed on particle size, and Dale and King have summarized the work of a number of experimentalists (124). They point out that it is well known that interstitial, intraperitoneal, and intratracheal injection of particulate silica produces only slow, chronic, silicotic fibrosis with no acute toxic reaction. On the other hand, both intratracheal and i.v. injections of finely divided silica of colloidal dimensions (20 A° or 2 μ) into rats (16 mg per 250 g rat) produced rapid

shock-like death. Others obtained similar results with other animals (also see Table 4 this monograph). From their survey they concluded that the larger-sized particles (ca 0.1 to 10 μ) produce the slow chronic fibrogenic response of silicosis and no acute toxic effect, and that molecularly dispersed silicic acid is freely diffusible through animal tissues and membranes and is nontoxic and readily excreted by the kidney. The intensely toxic range then involves "colloidal size" particles. These authors themselves investigated a variety of dusts of different composition and particle size, and observed, for example, that colloidal silica is more than 10 times as toxic as quartz of larger dimensions. A number of substances of particle size 0.1 to 10 μ appeared to be rather similar in the amounts necessary to kill, but there was an apparent break between their required amounts and those for the 20 A°-particle-size silica.

In studies on quartz 6 μ , scotch whinstone 6 μ , flint <1 μ , silicic acid (a soluble silica), and precipitated silica given orally (521, see section 5, Metabolism and Excretion) it was noted that there was a definite increase of urinary silica from the flint, the precipitated silica, and, to a greater extent, from the silicic acid. If one considers urinary excretion as an indication of absorption and circulation in the blood stream with the possibility of accumulation and hence of tissue toxicity, it would seem that there is a correlation between toxicity and particle size indicated in these results.

Another example of the effect of silica of large particle size may be found in the work of Bertke (061, section 1 Short Term

Studies). Diatomaceous earth, of particle size range 640-0.46 μ with 55% $<12 \mu$, when fed to rats 5% in their daily ration for 90 days caused no toxic symptoms. Nor was there any deviation from normal in histological sections or in silica content of the organs.

A pertinent study on particles of the colloidal range was that of Markovic and Arambasic (329) (section 3, Renal Mechanisms and Lesions) who induced chronic interstitial nephritis in guinea pigs by administering quartz of particle size 1-3 μ , 250 mg/liter in drinking water for several months. Lesions were formed that were quite similar to those found in man in Yugoslavia -- an endemic nephropathy caused by rock erosion in village communities on the banks of the lower reaches of large rivers.

Lundgren and Swensson (313) have pointed out, in regard to the pathogenesis of silicosis, that it is generally considered that quartz particles with a magnitude of 0.5 to 1 to 3 μ are of the greatest practical importance. They easily make their way down into the alveoli of the lungs and are retained there. Particles larger than 12 to 15 μ are to a large extent retained by the filtering apparatuses of the upper respiratory passages, and do not as frequently make their way down into the alveoli. Also, the fibrogenous effect of these larger particles seems less pronounced. To study the fibrogenesis of different particle sizes, the authors injected into guinea pigs in a single dose intraperitoneally somewhat coarse crystalline silica, 10 mg coarse particles of amorphous silica, and amorphous silica ca 61 μ , ca 14 μ . Large, coarse opaque

proliferative areas were seen in the peritoneum with cellular polymorphism and frequent infiltration of round cells. As particle size diminished, the proliferations of the peritoneum decreased in size.

For assessment of differences in toxicity between amorphous silicas of almost the same particle size, the work of Byers and Gage (084, Short Term Studies section 5) is of interest. The samples, from different suppliers, of particle size 19 μ , 20 μ (when fresh and 60 μ later), and 25 μ were administered intratracheally. As was pointed out earlier, silica was accumulated in the lungs and progressively disappeared, with some then appearing in the liver and kidneys. After 6 months, however, only the sample which showed evidence of aggregation (sample 2) was found in the latter organs. This sample had shown the greatest tendency to aggregate, when tested in a dust chamber; there was also greater difficulty in preparing aqueous suspensions.

A study considering surface area as well as particle size was that of Goldstein and Webster (201). These authors injected intratracheally into rats size-graded silica particles of <1 μ , 1 to 3 μ , and 2 to 5 μ ; each group had a surface area of 600 sq. cm/ml. They found that the degree of fibrosis was greater for the two large size fractions, and on the basis of statistical analysis was apparently related to the quantity (by weight) of the silica injected or the particle size rather than to the surface area.

In one of the very few long-term studies that are available, Gardner and Cummings (176) administered quartz of size 1 to 3 μ ,

and 6 to 12 μ , as well as aluminum oxide 1 to 3 μ , to rabbits by i.v. injection. (See Long Term Studies). After two to three years, histological examination showed that particles were segregated in different locations according to their size, the largest in the pulmonary capillaries, those of intermediate size in the spleen and hepatic lymph node, and the finest in the liver. The finest particles were the most active, while those of 10 to 12 μ were much less irritating, exciting a simple foreign body reaction which progressed little in three years. Fine aluminum oxide was merely phagocytosed and produced no fibrosis. This last study seems to indicate that the reaction, while correlated with particle size, is specific and chemical rather than purely physical in character (176).

Several authors have shown that polymerization and depolymerization of silicic acids may take place in various parts of the animal body. For example, Sauer, Laughland and Davidson (433, section 1, Guinea Pigs, Metabolism and Excretion) interpret their results as indicating that depolymerization of silica sol took place in the peritoneal cavity after oral administration. Arnold, Sasse and Strecker (041, Rats section 1, Metabolism and Excretion) after injecting i.p. aerosols of two different surface areas, and silicic acid gel, found that particles in the collecting tubules of the kidneys were essentially all the same size for the three different groups. From this they postulated polymerization in the distal nephron sections, a mechanism also proposed by others (041). Furthermore Settle and Sauer (445) postulated a critical concentration of 18 mg/100 ml of urine, below which soluble silica did

not polymerize.

Sauer et al. (433) found that solubility of the silica compound in the gastrointestinal tract is the limiting factor in its absorption. They also found that when guinea pigs were dosed with tetraethyl orthosilicate dissolved in 2.0 ml of ethanol, urinary excretion values were considerably higher than those for sodium silicate and silica sol mixed with the animals' food.

Different forms of silica of almost identical silica content, size distribution, and silica solubility were tested i.v. in mice, providing an example of the different results obtained according to chemical form (535). The fibrosing action in the liver was used as the criterion of measurement. It was found that tridymite was the most rapid in producing acellular collagenous silicotic nodules -- cristobalite was next, quartz third, and fused silica was the least fibrogenic. Silica particles were found initially in the spleen but with very little fibrosis. No fibrosis occurred in any other tissue. The authors proposed a mechanism possibly concerned with crystal structure and surface activity.

The study by Newberne and Wilson (372), already discussed in the sections on Short Term Studies and Metabolism and Excretion, serves as an example of the different results obtained from different silica salts. Sodium and magnesium silicates caused microscopic renal lesions in dogs, although no lesions were found in animals fed SiO_2 or aluminum silicate.

A theory proposed by Brieger and Gross (074) concerns the crystalline structure of silica necessary for fibrogenic

action. These authors studied coesite (particle size 0.14μ) and quartz (particle size 0.44μ); they gave intratracheal injections to rats and found that the fibrogenic potential of the two was of the same order. They concluded that the tetrahedral configuration of the SiO_2 molecule is a prerequisite for silicogenic action. This tetrahedral theory was substantiated by Strecker (470) in his study on the masking of SiO_2 surfaces. He prepared Aerosil surfaces which were masked with a number of agents, as illustrated in Table 29. When the Aerosil was administered to mice, these surface compounds delayed the acute toxic reaction, but not for long — as can be seen from the data in the table. He concluded that the reaction mechanism was connected with the free SiOH surface of solid particles constructed from SiO_2 tetrahedrons.

A final aspect to be considered when discussing the mechanism of toxicity is the action at the cellular level. Important works on the cytotoxic effects of silica compounds and the liberation of lysosomal enzymes have been published in England by Allison and Harington (216,012,010), and in Italy by Comolli (106). A detailed discussion of each is beyond the scope of this monograph, but attention is called to the concept, discussed by all of them and summarized so clearly by Allison (012). It is that silica particles are toxic because they are efficiently taken up by macrophages, and can then react relatively rapidly with the membranes surrounding the secondary lysosomes. The particles and lytic enzymes then can escape into the cytoplasm, causing general damage. Allison suggests that hydrogen bonding of silicic acid with lipid and protein constituents

of the membrane accounts for the induced permeability. Silica released from killed macrophages is as cytotoxic as the original preparation, and it is suggested that repeated cycles of macrophage killing in vivo leads to the mobilization of fibroblasts and fibrogenesis characterizing the disease silicosis.

Table 29 (470)

Masking of Aerosil Surfaces

Std. = Hours

Tg. = Days

Aerosil surface		Dose in mg	Death rate
sil. Oberfläche		Dosis: mg	Todesrate:
Si-OH		20	100% in 48 Std.
$\begin{array}{c} \text{H H} \\ \\ \text{Si-O-C-C-OH} \\ \\ \text{H H} \end{array}$	●	20	100% nach 48 Std.
$\begin{array}{c} \text{H H} \\ \\ \text{Si-O-C-C-H} \\ \\ \text{H H} \end{array}$	●	30 60	90% in 10 Tg. 100% in 3 Tg.
$\begin{array}{c} \text{H H H H} \\ \\ \text{Si-O-C-C-C-H} \\ \\ \text{H H H H} \end{array}$	●	30 60	~50% in 14 Tg.
$\begin{array}{c} \text{H H H H H H H H} \\ \\ \text{Si-O-C-C-C-C-C-C-H} \\ \\ \text{H H H H H H H H} \end{array}$	●	30 60	~30% in 14 Tg.
$\begin{array}{c} \text{H H H H} \\ \\ \text{Si-O-C-C-C-H} \\ \\ \text{H H H H} \end{array}$	◐	20 40	— 10% in 14 Tg.
	◑	20 40	30% in 3 Tg. 50% in 3 Tg. 80% in 14 Tg.
$\begin{array}{c} \text{CH}_3 \\ \\ \text{Si-O-Si-CH}_3 \\ \\ \text{CH}_3 \end{array}$	●	60	—
	◐	60	—
$\begin{array}{c} \text{OH} \\ \\ \text{Si-O-Al} \\ \\ \text{OH} \end{array}$	●	30 60	20% in 60 Std. 40% in 60 Std.

Ill. 1: Toxicity of aerosil with various surface coverings.
(Fatalities in mice with intraperitoneal application.)

VII. Consumer Exposure Information

The consumer may be exposed to silica from numerous natural and environmental sources, as discussed in the section on Chemical Information. The FDA has issued regulations for the use of calcium silicate (020,021) and sodium aluminosilicate (018,019) as anticaking agents; they specify a maximum usage of 2%, in salt for the former chemical and dried eggs for the latter. It has regulated silicon dioxide (016) to be used as an anticaking agent in vitamin products, 0.5%; in special dietary products, 0.75%; in spices, 2%; in meat-curing compounds, 2%; and in flavoring powders, 2%. A further regulation (025) extends the use of SiO_2 as an anticaking agent to any food at a maximum level of 2% when it has been demonstrated to have the anticaking effect; it is to be used at 2% in baby foods, however, only if NaCl, or NaCl substitutes, is a component.

The WHO/FAO monograph on "Toxicological Evaluation of Some Food Colors, Emulsifiers, Stabilizers, and Anti-Caking Agents and Certain Other Substances", 1969, (165), set no limit on Acceptable Daily Intakes (ADI) for the various forms of silicon dioxide, aluminum silicate, calcium silicate, magnesium silicate (including talc) and sodium aluminosilicate.

The NAS/NRC GRAS Survey Report lists the values shown in Table 30 for usage levels of four silica compounds in regular foods. Also, the Survey tabulates the possible daily intake per food category and total dietary level, based on the food consumption by total

Table 30

NAS/NRC GRAS Survey Report for Usage Levels of Silica Compounds in Foods

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# FIRMS REPORTING	*** USUAL USE *** WTD MEAN, %	*** MAXIMUM USE *** WTD MEAN, %
CALCIUM SILICATE NAS 0055	01 BAKED GOODS(R)	*	.15500	.15500
	03 OTHER GRAIN(R)	*	.05440	.05440
	04 FATS OILS(R)	*	.19200	.19200
	10 MEAT PRODS(R)	4	.00980	.01950
	11 POULTRY(R)	*	.00691	.01111
	13 FISH PRODS(R)	*	.01700	.01700
	16 SOFT CANDY(R)	*	.01024	.01883
	21 SOUPS(R)	4	.03163	.03187
	22 SNACK FOODS(R)	4	.12843	.57733
	23 BEV TYPE I(R)	*	.00070	.00200
	25 NUT PRODS(R)	*	.00964	.01055
	27 GRAVIES(R)	*	.03322	.06562
SILICA AEROGEL NAS 0174 FEMA 3506	28 IMIT DAIRY(R)	*	.49330	.49330
	48 SEAS FLAVRS(R)	*	.30167	.70094
	01 BAKED GOODS(R)	*	.03226	.04935
	02 BREAK CERLS(R)	*	.00850	.00850
	03 OTHER GRAIN(R)	*	.07500	.07500
	05 MILK PRODS(R)	*	.04997	.05297
	10 MEAT PRODS(R)	*	.00764	.01973
	11 POULTRY(R)	*	.01000	.05000
	13 FISH PRODS(R)	*	.01000	.05000
	21 SOUPS(R)	5	.01969	.02193
	22 SNACK FOODS(R)	*	.00100	.00500
	23 BEV TYPE I(R)	*	.04300	.04300
SODIUM ALUMINOSILICATE NAS 0178	27 GRAVIES(R)	*	.05045	.10050
	30 HARD CANDY(R)	*	.00750	.00750
	48 SEAS FLAVRS(R)	*	.40644	.40644
	01 BAKED GOODS(R)	*	.60039	1.39242
	02 BREAK CERLS(R)	*	.00015	.00045
	03 OTHER GRAIN(R)	*	.03600	.03600
	04 FATS OILS(R)	*	.04000	.06000
	05 MILK PRODS(R)	*	.03848	.03975
	07 FROZEN DAIRY(R)	*	.00299	.00498
	10 MEAT PRODS(R)	*	.00014	.00047
	11 POULTRY(R)	*	.01000	.05000
	13 FISH PRODS(R)	*	.00100	.00100
SODIUM ALUMINOSILICATE (CONT.) NAS 0178	15 CONDENSED RELISH(R)	*	.17217	.34435
	19 SWEET SAUCE(R)	*	.30675	.30675
	20 GELATIN PUD(R)	*	.03000	.07000
	21 SOUPS(R)	4	.41969	.42044
	22 SNACK FOODS(R)	*	.13145	.92768
	23 BEV TYPE I(R)	*	.07919	.07919
	27 GRAVIES(R)	*	.06434	.19911
	28 IMIT DAIRY(R)	5	.87533	.87533
	48 SEAS FLAVRS(R)	*	.53930	.53950

sample, as shown in Table 31. In summary, the GRAS survey reports a high possible daily intake of calcium silicate for the 2-65 year old age group as ca 481 mg; for silica aerogel ca 340 mg, for sodium aluminum silicate ca 2278 mg and for sodium calcium aluminosilicate hydrate 2.66 mg.

The annual poundage data for calcium silicate, magnesium silicate and silica aerogel is reported in Table 32.

Table 31

Possible Daily Intakes of NAS Appendix A Substances (Groups I & II), Per Food Category and Total Dietary, Based on Food Consumption by Total Sample --- See Explanatory Notes in Exhibits Section

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# FIRMS REPORTING	*** USUAL USE *** WTD MEAN, %	*** MAXIMUM USE *** WTD MEAN, %
CALCIUM SILICATE NAS 0055	01 BAKED GOODS(R)	*	.15500	.15500
	03 OTHER GRAIN(R)	*	.05440	.05440
	04 FATS OILS(R)	*	.19200	.19200
	10 MEAT PRODS(R)	4	.00980	.01950
	11 POULTRY(R)	*	.00691	.01111
	13 FISH PRODS(R)	*	.01700	.01700
	16 SFT CANDY(R)	*	.01024	.01883
	21 SOUPS(R)	4	.03163	.03187
	22 SNACK FOODS(R)	4	.12843	.57733
	23 BEV TYPE I(R)	*	.00070	.00200
	25 NUT PRODS(R)	*	.00564	.01055
	27 GRAVIES(R)	*	.03322	.06562
	28 IMIT DAIRY(R)	*	.49330	.49330
	48 SEAS FLAVRS(R)	*	.30167	.70094
SILICA AEROGEL NAS 0174 FEMA 3506	01 BAKED GOODS(R)	*	.03226	.04935
	02 BREAK CERLS(R)	*	.00850	.00350
	03 OTHER GRAIN(R)	*	.07500	.07500
	05 MILK PRODS(R)	*	.04997	.05297
	10 MEAT PRODS(R)	*	.00764	.01973
	11 POULTRY(R)	*	.01000	.05000
	13 FISH PRODS(R)	*	.01000	.05000
	21 SOUPS(R)	5	.01959	.02183
	22 SNACK FOODS(R)	*	.00100	.00500
	23 BEV TYPE I(R)	*	.04300	.04300
	27 GRAVIES(R)	*	.05045	.10050
	30 HARD CANDY(R)	*	.00750	.00750
	48 SEAS FLAVRS(R)	*	.40644	.40644
SODIUM ALUMINOSILICATE NAS 0178	01 BAKED GOODS(R)	*	.68039	1.39242
	02 BREAK CERLS(R)	*	.00015	.00045
	03 OTHER GRAIN(R)	*	.03600	.03600
	04 FATS OILS(R)	*	.04000	.06000
	05 MILK PRODS(R)	*	.03848	.03975
	07 FROZEN DAIRY(R)	*	.00299	.00498
	10 MEAT PRODS(R)	*	.00014	.00047
	11 POULTRY(R)	*	.01000	.05000
	13 FISH PRODS(R)	*	.00100	.00100
	15 CONCN RELSH(R)	*	.17217	.34435
SODIUM ALUMINOSILICATE (CONT.) NAS 0178	19 SWEET SAUCE(R)	*	.38675	.38675
	20 GELATIN PUD(R)	*	.03000	.07000
	21 SOUPS(R)	4	.41969	.42044
	22 SNACK FOODS(R)	*	.63145	.92768
	23 BEV TYPE I(R)	*	.07918	.07918
	27 GRAVIES(R)	*	.06434	.19911
	28 IMIT DAIRY(R)	5	.87533	.87533
	48 SEAS FLAVRS(R)	*	.53938	.53950

Table 31

Possible Daily Intakes of NAS Appendix A Substances (Groups I & II). Per Food Category and Total Dietary, Based on Food Consumption by Total Sample --- See Explanatory Notes in Exhibits Section

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# OF FIRMS	AGE	***** AVERAGE	POSSIBLE DAILY INTAKE, MG. HIGH A	***** HIGH B
CALCIUM SILICATE NAS 0055	01 BAKED GOODS(R)	*	0-5 MO.	5.270000	6.975000	5.270000
			6-11 MO.	39.370000	80.290000	39.370000
			12-23 MO.	84.475000	139.190000	84.475000
			2-65+ YR.	212.660000	315.890000	212.660000
CALCIUM SILICATE NAS 0055	03 OTHER GRAIN(R)	*	0-5 MO.	.272000	.924800	.272000
			6-11 MO.	5.276000	15.558400	5.276800
			12-23 MO.	8.921600	26.617600	8.921600
			2-65+ YR.	15.123200	33.401600	15.123200
CALCIUM SILICATE NAS 0055	04 FATS OILS(R)	*	0-5 MO.	.960000	.960000	.960000
			6-11 MO.	5.376000	14.400000	5.376000
			12-23 MO.	12.096000	23.040000	12.096000
			2-65+ YR.	33.600000	60.672000	33.600000
CALCIUM SILICATE NAS 0055	10 MEAT PRODS(R)	4	0-5 MO.	.107800	.284200	.214500
			6-11 MO.	2.028600	5.468400	4.036500
			12-23 MO.	2.959600	5.086200	5.889000
			2-65+ YR.	7.683200	12.749800	15.288000
CALCIUM SILICATE NAS 0055	11 POULTRY(R)	*	0-5 MO.	.034550	.158930	.055550
			6-11 MO.	.269490	.912120	.433290
			12-23 MO.	.456060	1.271440	.733260
			2-65+ YR.	.891390	2.266480	1.433190
CALCIUM SILICATE NAS 0055	13 FISH PRODS(R)	*	0-5 MO.	.017000	.051000	.017000
			6-11 MO.	.221000	.833000	.221000
			12-23 MO.	.918000	2.255000	.918000
			2-65+ YR.	2.108000	5.253000	2.108000
CALCIUM SILICATE NAS 0055	16 SOFT CANDY(R)	*	0-5 MO.	.020480	.204800	.037660
			6-11 MO.	.225280	.696320	.414260
			12-23 MO.	.358400	.952320	.659050
			2-65+ YR.	.593520	1.802240	1.092140
CALCIUM SILICATE NAS 0055	21 SCUPS(R)	4	0-5 MO.	.063260	.474450	.063740
			6-11 MO.	7.365790	22.995010	7.425710
			12-23 MO.	11.007240	30.396430	11.050760
			2-65+ YR.	10.026710	26.727350	10.102790
CALCIUM SILICATE NAS 0055	22 SNACK FOODS(R)	4	0-5 MO.	*****	.128430	*****
			6-11 MO.	.513720	1.412730	2.309320
			12-23 MO.	1.412730	3.981330	6.350630
			2-65+ YR.	1.669550	4.751910	7.505290
CALCIUM SILICATE NAS 0055	23 BEV TYPE I(R)	*	0-5 MO.	.016800	.025200	.048000
			6-11 MO.	.158500	.543900	.454000
			12-23 MO.	.379400	1.137500	1.084000
			2-65+ YR.	.728000	1.943900	2.080000

Table 31

Possible Daily Intakes of NAS Appendix A Substances (Groups I & II), Per Food Category and Total Dietary, Based on Food Consumption by Total Sample --- See Explanatory Notes in Exhibits Section

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# OF FIRMS	***** (AGE)	***** AVERAGE	***** HIGH A	***** HIGH B
CALCIUM SILICATE NAS 0055	25 NUT PRODS(R)	*	C-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	***** .208660 .152280 .293200	.011280 .755760 .567600 .874200	***** .390350 .264850 .548600
CALCIUM SILICATE NAS 0055	27 GRAVIES(R)	*	C-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.033270 .465080 1.195920 2.757260	.099660 1.295500 3.388440 7.075860	.065620 .918680 2.362320 5.446460
CALCIUM SILICATE NAS 0055	28 IMIT DAIRY(R)	*	C-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.000000 6.906200 3.946400 4.439700	.000000 11.345900 16.772200 7.399500	.000000 6.906200 3.946400 4.439700
CALCIUM SILICATE NAS 0055	43 SEAS FLAVRS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	***** ***** ***** .030167	***** ***** ***** .060334	***** ***** ***** .070094
CALCIUM SILICATE NAS 0055	ALL CATEGORIES ***** ***** *****	13	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	6.795110 68.389540 128.278630 292.604417	10.297750 156.537287 248.696394 480.958675	7.004070 73.532110 138.810870 311.497464
SILICA AEROGEL NAS 0174	01 BAKED GOODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	1.096840 8.194040 17.581700 44.260720	1.451700 16.710680 28.969480 65.745880	1.677900 12.534900 26.895750 67.708200
SILICA AEROGEL NAS 0174	02 .BREAK CERLS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.051000 1.895500 2.218500 1.700000	.144500 5.083000 4.326500 4.403000	.051000 1.895500 2.218500 1.700000
SILICA AEROGEL NAS 0174	03 OTHER GRAIN(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.375000 7.275000 12.300000 20.850000	1.275000 21.450000 28.425000 46.050000	.375000 7.275000 12.300000 20.850000
SILICA AEROGEL NAS 0174	05 MILK PRODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	2.698380 31.181280 27.233050 19.738150	1.998800 149.959970 87.147680 60.263820	2.860380 33.053280 28.868650 20.923150

Table 31

Possible Daily Intakes of NAS Appendix A Substances (Groups I & II), Per Food Category and Total Dietary, Based on Food Consumption by Total Sample --- See Explanatory Notes in Exhibits Section

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# OF FIRMS	***** (AGE)	Possible Daily Intake, MG. AVERAGE	HIGH A	HIGH B
SILICA AEROGEL NAS 0174	10 MEAT PRDGS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.004040 1.581480 2.307280 5.989760	.221560 4.263120 3.965160 9.939640	.217030 4.084110 5.958460 15.468320
SILICA AEROGEL NAS 0174	11 POULTRY(R)	*	0-5 MO. 6-11 MO. 12-23 MO.	.050000 .390000 .660000	.230000 1.320000 1.840000	.250000 1.950000 3.300000
SILICA AEROGEL NAS 0174	13 FISH PRDGS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.010000 .130000 .540000 1.240000	.030000 .490000 1.350000 3.090000	.050000 .650000 2.700000 6.200000
SILICA AEROGEL NAS 0174	21 SOUPS(R)	5	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.039300 4.507770 6.852120 6.241730	.295350 14.314630 18.922090 16.638050	.043660 5.086390 7.558840 6.920110
SILICA AEROGEL NAS 0174	22 SNACK FOODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	***** .004000 .011000 .013000	.001000 .011000 .031000 .037000	***** .020000 .055000 .065000
SILICA AEROGEL NAS 0174	23 BEV TYPE I(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	1.032000 9.761000 23.306000 44.720000	1.540000 33.411000 69.874000 119.411000	1.032000 9.761000 23.306000 44.720000
SILICA AEROGEL NAS 0174	27 GRAVIES(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.050450 .706300 1.816200 4.187350	.151350 1.967550 5.145900 10.745850	.100500 1.407000 3.518000 8.341500
SILICA AEROGEL NAS 0174	30 HARD CANDY(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.000000 .007500 .022500 .045000	.000000 .027500 .067500 .127500	.000000 .007500 .027500 .045000
SILICA AEROGEL NAS 0174	48 SEAS FLAVRS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	***** ***** ***** .040644	***** ***** ***** .081288	***** ***** ***** .040644
SILICA AEROGEL NAS 0174	ALL CATEGORIES	16	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	5.487090 65.713070 94.648950 150.316354	7.347260 249.044094 250.146598 335.934960	6.557470 77.724680 116.839700 159.431924

Table 31

Possible Daily Intakes of NAS Appendix A Substances (Groups I & II). Per Food Category and Total Dietary, Based on Food Consumption by Total Sample --- See Explanatory Notes in Exhibits Section

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# OF FIRMS	***** (AGE)	***** AVERAGE	***** HIGH A	***** HIGH B
SODIUM ALUMINOSILICATE NAS 0175	01. BAKED GOODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	23.133260 172.819060 370.812550 933.495080	30.617550 352.442020 610.990220 1386.634820	47.342260 353.674680 758.868900 1910.400240
SODIUM ALUMINOSILICATE NAS 0176	02. BREAK CERLS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.000900 .033450 .039150 .030000	.002550 .089700 .076350 .077700	.002700 .100350 .117450 .090000
SODIUM ALUMINOSILICATE NAS 0178	03. OTHER GRAIN(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.180000 3.492000 5.904000 10.008000	.612000 10.256000 13.644000 22.104000	.180000 3.492000 5.904000 10.008000
SODIUM ALUMINOSILICATE NAS 0176	04. FATS OILS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.200000 1.120000 2.520000 7.000000	.200000 3.000000 4.900000 12.440000	.300000 1.600000 3.700000 10.500000
SODIUM ALUMINOSILICATE NAS 0178	05. MILK PRODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	2.077920 24.011520 20.971600 15.195600	1.539200 115.478480 67.109120 46.406880	2.145500 24.804000 21.663750 15.701250
SODIUM ALUMINOSILICATE NAS 0178	07. FROZEN DAIRY(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.029900 .284050 .430560 .765440	.122590 .785360 1.010620 1.844830	.045800 .473100 .717120 1.274880
SODIUM ALUMINOSILICATE NAS 0178	10. MEAT PRODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.001540 .028980 .042280 .109760	.004060 .078120 .072660 .182140	.005170 .097250 .141940 .366480
SODIUM ALUMINOSILICATE NAS 0178	11. POULTRY(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.090000 .390000 .660000 1.250000	.230000 1.320000 1.840000 3.280000	.250000 1.500000 3.300000 6.450000
SODIUM ALUMINOSILICATE NAS 0178	13. FISH PRODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.001000 .013000 .054000 .124000	.003000 .040000 .135000 .305000	.001000 .013000 .054000 .124000
SODIUM ALUMINOSILICATE NAS 0178	15. CANNED RELISH(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	***** 1.377360 4.820750 15.150960	.177170 3.797740 13.084920 36.500040	***** 2.754800 9.641800 30.302800

Table 31

Possible Daily Intakes of NAS Appendix A Substances (Groups I & II). Per Food Category and Total Dietary, Based on Food Consumption by Total Sample --- See Explanatory Notes in Exhibits Section

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# OF FIRMS	***** (AGE)	***** AVERAGE	***** HIGH A	***** HIGH B
SODIUM ALUMINOSILICATE NAS 0178	19 SWEET SAUCE(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	1.160250 3.480750 10.055000 26.299000	1.547000 11.989250 29.393000 69.225250	1.180250 3.480750 10.055000 26.299000
SODIUM ALUMINOSILICATE NAS 0178	20 GELATIN PUD(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.600000 3.840000 4.140000 6.120000	.810000 11.640000 10.080000 15.750000	1.400000 8.960000 9.660000 14.280000
SODIUM ALUMINOSILICATE NAS 0178	21 SCUPS(R)	4	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.839380 97.787770 146.052120 123.041730	6.295350 305.114630 403.222090 354.638050	.840880 97.982520 146.313120 133.279480
SODIUM ALUMINOSILICATE NAS 0178	22 SNACK FOODS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	***** 2.525800 6.945950 8.202850	.631450 6.945950 19.574950 23.363650	***** 3.710720 10.204480 12.059840
SODIUM ALUMINOSILICATE NAS 0178	23 BEV TYPE I(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	1.900320 17.973860 42.915560 82.347200	2.650480 61.522860 128.667500 219.882860	1.900320 17.973860 42.915560 82.347200
SODIUM ALUMINOSILICATE NAS 0178	27 GRAVIES(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.064340 .900760 2.316240 5.340220	.193020 2.509260 6.562660 13.704420	.199110 2.787540 7.167960 16.526130
SODIUM ALUMINOSILICATE NAS 0178	28 IPIT DAIRY(R)	5	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	.000000 12.254620 7.002640 7.877970	.000000 20.132590 29.761220 13.129550	.000000 12.254620 7.002640 7.877970
SODIUM ALUMINOSILICATE NAS 0178	48 SEAS FLAVRS(R)	*	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	***** ***** ***** .053938	***** .053938 .107876 .267650	***** ***** ***** .053950
SODIUM ALUMINOSILICATE NAS 0178	ALL CATEGORIES ***** ***** *****	20	0-5 MO. 6-11 MO. 12-23 MO. 2-65+ YR.	30.200810 347.332980 625.632910 1252.461748	45.830420 907.230890 1340.232206 2219.946280	55.770010 536.166230 1037.506220 2277.943220

Table 31

Possible Daily Intakes of NAS Appendix A Substances (Groups I & II). Per Food Category and Total Dietary. Based on Food Consumption by Total Sample --- See Explanatory Notes in Exhibits Section

SUBSTANCE NAME (SURVEY NO.)	FOOD CATEGORY NO. NAME	# OF FIRMS	***** (AGE)	POSSIBLE DAILY INTAKE, MG. *****		
				AVERAGE	HIGH A	HIGH B
SODIUM CAL ALUMSIL HYD NAS 0184	10 MEAT PRDCE(R)	*	0-5 MO.	.011000	.029000	.055000
			6-11 MO.	.207000	.558000	1.025000
			12-23 MO.	.302000	.519000	1.510000
			2-65+ YR.	.784000	1.301000	3.920000
SODIUM CAL ALUMSIL HYD NAS 0184	11 POULTRY(R)	*	0-5 MO.	.005000	.023000	.025000
			6-11 MO.	.039000	.132000	.195000
			12-23 MO.	.066000	.184000	.330000
			2-65+ YR.	.129000	.328000	.645000
SODIUM CAL ALUMSIL HYD NAS 0184	21 SEUPS (R)	*	0-5 MO.	.002000	.015000	.010000
			6-11 MO.	.233000	.727000	1.165000
			12-23 MO.	.348000	.961000	1.740000
			2-65+ YR.	.317000	.845000	1.585000
SODIUM CAL ALUMSIL HYD NAS 0184	22 SNACK FOODS(R)	*	0-5 MO.	*****	.005000	*****
			6-11 MO.	.020000	.055000	.040000
			12-23 MO.	.055000	.155000	.110000
			2-65+ YR.	.065000	.185000	.130000
SODIUM CAL ALUMSIL HYD NAS 0184	ALL CATEGORIES ***** ***** *****	*	0-5 MO.	.018000	.072000	.090000
			6-11 MO.	.499000	1.472000	2.435000
			12-23 MO.	.771000	1.819000	3.690000
			2-65+ YR.	1.295000	2.659000	6.280000

Table 32 Animal Poundage Data for NAS Appendix A

	Reports to NAS 1960/70	1960	1970	Total 1970 Poundage to NAS	Poundage to FEMA	NAS + FEMA
Calcium Silicate	13/15	1,012,650	1,154,391	1,290,466		1,290,466
Magnesium Silicate	5/5	52,175	47,144	47,144		47,144
Silica Aerogel	14/16	66	23,277	29,627	6,666	36,293

- 1 Achenbach, K. 1968
Turbidity reduction
U.S. Pat. 3,413,120 issued in 1968
- 2 Achenbach, K. 1968
Verfahren zur entfernung von kolloidale trubungen
bidenden eiweisstoffen aus garungsgetranken,
insbesondere aus bier. (Method for removing
colloidal haze-forming proteins from fermented
beverages, particularly from beer)
W. Ger. Pat. 1,442,338 issued in 1968
- 3 Afanas'eva, L.V. 1953
Determination of silica in blood and organs
photocolorimetrically
Biokhimiya 18:319-323
- 4 Afremov, Leonard C., and John T. Vandenberg. 1966
High-resolution spectra of inorganic pigments in
the mid-infrared region from 1500 cm.⁻¹ to 200
cm.⁻¹
J. Paint Technol. 38(495):169-202
- 5 Akazaki, Kaneyoshi, and Yozo Inagaki. 1959
Experimental anthracosis, anthracosilicosis and
the relation of these to tuberculosis in
complication
Tohoku J. Exp. Med. 71:195-205
- * 6 Akiya, S., T. Misawa, N. Motohashi, Y. Nakazawa,
T. Osawa, A. Tanimura, T. Niwaguchi, and T.
Niwase. 1959
Studies on the silica in animal tissue. The
relationship between silica in drinks and foods
and in human blood vessels
Bull. Tokyo Med. Dental Univ. 6(3):393-411
- 7 Alberti, Carlo. 1939
Untersuchung uber capsicum aus Italienisch
Somaliland (Capsicum frutescens L.)
Ann. Chim. Appl. 29:392-402
- 8 Aleksandrov, S.E. 1963
On lesions of abdominal organs caused by talc
Zdravookhr. Beloruss. 9:34-36
- 9 Alexanian, C., P. Morel, and L. Le Bouffant. 1966
Infrared absorption spectra of natural minerals
Bull. Soc. Fr. Ceram. 71:3-38
- * 10 Allison, A. 1970
Effects of particles on lysosomes
Adv. Sci. 27:137-140
- 11 Allison, A.C., and P. D'Arcy Hart. 1968
Potentiation by silica of the growth of
Mycobacterium tuberculosis in macrophage cultures
Brit. J. Exp. Pathol. 49(5):465-476
- * 12 Allison, A.C., J.S. Harington, and M. Birbeck.
1966
The cytotoxic effects of silica on macrophages
J. Exp. Med. 124(2):141-154
- 13 Andersson, Lars Henrik. 1962
Studies in the spectrometric and gravimetric
determination of silica
Ark. Kemi 19(21):257-264
- 14 Ando, T., M. Kamoda, F. Onda, H. Ito, T.
Shirasaki, and T. Miki. 1968
Occlusion of filtration-impeding substances in
sugar crystal
Proc. Int. Soc. Sugar Cane Technol. 13:429-434
- 15 Ando, T., M. Kamoda, F. Onda, H. Ito, T.
Shirasaki, and T. Miki. 1969
Occlusion of filtration-impeding substances in
sugar crystals
Proc. Res. Soc. Jap. Sugar Refineries Technol.
1969(21):112-116
- * 16 Anon. 1962
Food additives. Silicon dioxide
Fed. Regist. 27:4914
- 17 Anon. 1963
Food additives. Resinous and polymeric coatings
Fed. Regist. 28:8024
- * 18 Anon. 1963
Dried eggs, dried whole eggs; order amending
standard of identity
Fed. Regist. 28:7776
- * 19 Anon. 1963
Dried egg yolks; standard of identity to provide
for optional anticaking ingredient sodium
silicoaluminate
Fed. Regist. 28:4179-4180
- * 20 Anon. 1963
Food additives. Calcium silicate
Fed. Regist. 28:12747-12748
- * 21 Anon. 1963
Food additives. Calcium silicate
Fed. Regist. 28:7559
- 22 Anon. 1964
Food additives. Silicon dioxide
Fed. Regist. 29:14224
- 23 Anon. 1964
Food additives. Defoaming agents
Fed. Regist. 29:2780-2781
- 24 Anon. 1964
Monographs for a new German pharmacopeia
Pharm. Ind. 26(4):223-226
- * 25 Anon. 1966
Food additives. Silicon dioxide
Fed. Regist. 31:9106
- 26 Anon. 1968
Food additives. Microcapsules for flavoring oils
Fed. Regist. 33(232):17752
- 27 Anon. 1968
Single-pass filtration for beer
Process Biochem. 3(9):39
- 28 Anon. 1970
Photometric determination of residual silicic acid
Silikat-Technik 21(5):173-174
- 29 Anon. 1970
Beer stabilization
Int. Brewers' J. 106(1254):39-41
- 30 Anon. 1970
Der drehfilter - neu in der kellerpraxis. (The
rotary filter - the latest in cellar practice)
Alkohol-Industrie 83(13):311-313
- 31 Anon. 1971
The silicates in food
Brit. Food J. 73(842):80-81
- 32 Anon. 1971
Navel ills
Brit. Med. J. 1:248
- 33 Antopol, W. 1933
Lycopodium granuloma; its clinical and pathologic
significance, together with note on granuloma
produced by talc
Arch. Path. 16:326-331
- 34 Appleton, Martin D., and Richard J. Burns. 1958
A modification of the colorimetric analysis for
silica according to the method of Gettler and
Umberger
Proc. Pa. Acad. Sci. 32:69-73
- 35 Ardelean, I. 1967
Action of silicium dioxide, caol and calcium
carbonate powders on immunity reactions
Fiziol. Norm. Pat. 13:385-390
- 36 Ardelean, I. 1968
Action of silicon dioxide, coal and calcium
carbonate dust on immunity responses
Rom. Med. Rev. 12:3-7

- 37 Ardoino, L.A., V. Caputo, and M. Greco. 1961
Protein modifications induced by silica in guinea
pig serum in relation to decrease in complement
titer
Med. Lav. 52:601-604
- 38 Arkhipenko, D.K., and E.A. Plekhanova. 1961
Infrared spectra of kaolinite and products of its
degradation
Izv. Sib. Otd. Akad. Nauk S.S.S.R.
1961(3):109-112
- 39 Armour & Co. 1966
Antacids
Ger. Pat. 1,220,554 issued Jul. 7, 1966
- 40 Arnaud, M. 1932
Treatment by injections of colloidal suspensions
of silica and calcium oxide
Rev. Tech. Chir. 24:61-64
- * 41 Arnold, Michael, Dieter Sasse, and Franz Josef
Strecker. 1961
Behavior of Aerosil in the animal body
Beitr. Silikose-Forsch., Sonderb. 4:87-97
- 42 Aschaffenburg Zellstoffwerke Akt.-Ges. (by
Konrad Bernhauer, and Wilhelm Friedrich). 1958
Recovery of B12 vitamins in concentrated form
Ger. Pat. 1,037,066 issued Aug. 21, 1958
- * 43 AtLee, W.E., Jr. 1972
Talc and cornstarch emboli in eyes of drug abusers
J. Am. Med. Assoc. 219:49-51
- 44 Aumonier, P., and R. Quilichini. 1962
Photometric estimation of silicon in the presence
of phosphorus by the formation of reduced
molybdosilicous acid. Application to blood serum
Ann. Biol. Chim. (Paris) 20(10-12):965-971
- * 45 Autio, Lauri, and Martti Turunen. 1966
The effect of intravenous silica on serum
electrophoresis in dog
Ann. Med. Exp. Biol. Fenn. 42(3):132-135
- 46 Autio, Lauri, Martti Turunen, and Arto Lahtiharju.
1966
Experimental silica cirrhosis in dog
Ann. Med. Exp. Biol. Fenn. 42(4):173-176
- 47 Avy, Alban. 1955
Determination of free silica in the presence of
combined silica
Mem. Serv. Chim. Etat. (Paris) 40:345-362
- 48 Baba, Kenichi, and Akira Baba. 1959
Relation between protein and silicate in rice
plants (Jap.)
Nippon Nogei Kagaku Kaishi 33(8):710-714
- 49 Bachrach, E., and M. Lefevre. 1929
Importance of silica in living organisms; biology
of diatoms
J. Physiol. Pathol. Gen. 27:241-249
- 50 Bailey, C.B. 1967
Silica excretion in cattle fed a ration
predisposing to silica urolithiasis:
Total excretion and diurnal variations
Am. J. Vet. Res. 28(127):1743-1749
- 51 Bailey, C.B. 1967
Siliceous urinary calculi in calves: prevention
by addition of sodium chloride to the diet
Science 155:696-697
- 52 Baker, Joe H. 1966
Cattle feeding
U.S. Pat. 3,244,527 issued Apr. 5, 1966
- 53 Baker, Robert H., and Charles Barkenbus. 1937
Analyse von whiskygrunspan
Ind. Eng. Chem. 29:755-757
- 54 Barrett, George S. 1951
Influence of alumina gels on prevention of
urinary calculi
J. Urol. 66:315-330
- 55 Bass, Arnold M., and Herbert P. Broida. 1957
Sharp line absorption in silicon near 3400 cm.-1
J. Opt. Soc. Am. 47:163-164
- * 56 Beamer, W.D., M.H.P. Friedman, and P.T. Chu. 1953
Influence of cortisone on formation of
intraperitoneal adhesions
Exp. Med. Surg. 11(4):241-247
- 57 Beekman, Stewart M. 1961
Darstellung und eigenschaften neuer, gastrischer
antacida. 3. Mitt. Aluminiumhydroxid-glycin
J. Pharm. Sci. 50:571-577
- 58 Beekman, Stewart M. 1961
Darstellung und eigenschaften neuer gastrischer
antacida. 4. Mitt. Aluminiumhydroxid-
magnesiumtrisilicat
J. Pharm. Sci. 50:577-580
- 59 Benecke, E. 1933
Reaction of subcutaneous and peritoneal
connective tissue to injection of stone dust
containing silicon
Beitr. Z. Path. Anat. Z. Allg. Path. 91:503-514
- 60 Benvegnin, L., and E. Capt. 1950
La filtration des vins sur terres de diatomées.
(The use of silicious earth in the filtration of
wines)
Rev. Romande Agric. Vitic. Arboric. 6(7):52-54
- * 61 Bertke, Eldridge M. 1964
The effect of ingestion of diatomaceous earth in
white rats: A subacute toxicity test
Toxicol. Appl. Pharmacol. 6(3):284-291
- 62 Bezeau, L.M., A. Johnston, and S. Smoliak. 1966
Silica and protein content of mixed prairie and
fescue grassland vegetation and
its relationship to the incidence of silica
urolithiasis
Can. J. Plant Sci. 46(6):625-631
- 63 Bezeau, L.M., C.B. Bailey, and S.B. Slen. 1961
Silica urolithiasis in beef cattle. IV. The
relationship between the pH and buffering
capacity of the ash of certain feeds, pH of the
urine, and urolithiasis
Can. J. Anim. Sci. 41(1):49-54
- 64 Bhatt, M.P., and Dinkar S. Datar. 1968
Caking of salt
Salt Res. Ind. 5(3/4):57-67
- 65 Bibus, B. 1966
Talcum granuloma in the area of the ureter
Z. Urol. 59:525-528
- 66 Bigger, Joseph W., and J. Havelock Nelson. 1941
The growth of coliform bacilli in distilled water
J. Pathol. Bact. 53(2):189-206
- 67 Bloomfield, J.J., R.R. Sayers, and P.H. Goldman.
1935
Die ausscheidung von kieselstaure im urin durch
kieselstaub ausgesetzten personen
Public Health Rep. 50:421-424
- 68 Bobovich, Ya. S., and T.P. Tulub. 1960
Temperature dependence of combination scattering
band intensity in crystalline and glassy bodies.
I. Experimental investigation in the Stokes
spectral region
Opt. Spektrosk. 9:747-753
- 69 Boqdan, I.G., T.F. Gorb, and M.M. Polyachenko.
1969
Use of diatomaceous earth in the fats-and-oils
industry (Russ.)
Pishch. Prom. (Kiev) 1969(10):38-40
- 70 Brachetto-Brian, Domingo. 1926
Accion de las inoculaciones de kieselgur en el
tejido muscular de la gallina. (Reaction in
fowls to intramuscular injections of infusorial
earth (kieselgur))
Rev. Soc. Argent. Biol. 2(6):431-438

- 71 Brain, Domingo Brachetto. 1926
Accion de las inoculaciones de kieselguhr en el tejido muscular de la gallina. (Results of inoculations of infusorial earth into the muscular tissues of fowls)
Sem. Med. 33(1720):1752-1755
- 72 Brandt, R., and H. Plotnick. 1956
Multiple cutaneous and subcutaneous sarcoid-like foreign body granulomas: case with parallel course in various sites
Am. Med. Assoc. Arch. Dermatol. 74:128-130
- 73 Brenner, P. 1968
Schichtenfiltration und filterschichten. (Sheet filtration and filter sheets)
Brauwelt 108(64):1161-1165
- * 74 Brieger, H. 1966
On the theory of silicosis. I. Coesite
Arch. Environ. Health (Chicago) 13:38-43
- 75 Brock, Joachim. 1937
Calcinosis circumscripta with sclerodactylism in infancy
Z. Kinderheilkd. 58:751-756
- 76 Brody, J. 1970
Ingredient water: analytical requirements
Food Eng. 42(5):94-95
- 77 Brown, Ernest A. 1957
The adsorption of lipids from the erythrocyte surface by silica and alumina
J. Cell. Comp. Physiol. 50(1):49-56
- 78 Bruger, Maurice. 1935
The state of cholesterol and the nature of the cholesterol-protein complex in pathological body fluids
J. Biol. Chem. 108(2):463-470
- 79 Brune, Heinrich. 1952
The influence of adsorptive materials on resorption
Arch. Tierernahr. 3:1-31
- 80 Bsteh, O. 1966
Clinical manifestations of talcum granulomatosis of the peritoneum
Int. Surg. 45:424-428
- 81 Bundus, R.H., and P.P. Noznick. 1970
Powdered vinegar
Can. Pat. 847,288 issued in 1970
- 82 Burness, A.T. 1967
Use of kieselguhr to increase cell production for animal virus investigations
Nature (London) 213:833
- * 83 Butz, W.C. 1970
Disseminated magnesium and aluminum silicate associated with paregoric addiction
J. Forensic Sci. 15:581-587
- * 84 Byers, P.D., and J.C. Gage. 1961
The toxicity of precipitated silica
Brit. J. Ind. Med. 18:295-302
- 85 Campbell, J.A. 1931
Effects of precipitated silica and of iron oxide on incidence of primary lung tumors in mice
Brit. Med. J. 2:275-280
- 86 Campen, W.A.C. 1952
Simple and rapid determination of silica in cattle feeds
Chem. Weekbl. 48:579
- 87 Carminati, V., and M. Chirico. 1952
Tumor formation by silicic anhydride in rats treated contralaterally with 3,4-benzopyrene
Atti Soc. Lomb. Sci. Med. Biol. 7:400-401
- 88 Carminati, Valentino. 1931
Ricerche sull'azione neofornativa della farina fossile in combinazione con altri stimoli e sugli inesti seriati del tessuto di granulazione. (Neofornative action of kieselguhr in combination with other stimuli; and on serial grafts of granulation tissue)
Tumori 17(2):101-127
- 89 Carminati, Valentino, and Mario Chirico. 1951
Action on the connective tissue of some substances contained by kieselguhr.
II. Strontium sulfate and calcium carbonate
Tumori 25:198-207
- 90 Carminati, Valentino, and Mario Chirico. 1951
Action of some kieselguhr substances on the connective tissue. I. Silica
Tumori 27:83-92
- 91 Carter, S.B. 1965
Problems in interpreting teratogenic effects in eggs
Proc. Eur. Soc. Stud. Drug Toxicity 5:142-148
- 92 Castro, L. De P., and H.P. Resende. 1969
Efeito, in vivo, de alguns antiacidos e alimentos sobre o pH do conteúdo gastrico. Sua importancia na terapeutica da ulcer peptica. (In vivo effect of some antiacids and foods on the pH of the gastric content. Its importance in the treatment of peptic ulcer)
Rev. Assoc. Med. Minas. Gerais. 19(1):53-61
- 93 Catalano, F.E. 1946
Granulomas due to talc used on surgical gloves; experimental study
Dia Med. 18:1582-1588
- 94 Cerix Puffed Wheat Co., Concord. 1929
Tapiocapraparat
Austral. Pat. 17,248/1928 issued Dec. 31, 1929
- 95 Cesar, Leyton G. 1945
Chemical determination of silica in biological products
An. Quim. Farm. 1945(35):1-3
- 96 Chang, Ching-Ming, Chin-Chih Liu, and Shu-Hsien Wang. 1963
Rapid determination of SiO₂ by precipitation as potassium fluorosilicate
Kuei Suan Yen Hsueh Pao 2(3):163-171
- 97 Chapman, John S. 1963
Eosinophil-stimulating properties of certain chemical substances
Am. J. Clin. Pathol. 40(4):357-362
- 98 Chatterjee, Anil C. 1968
Chemicals used in white sugar manufacture
Indian Sugar 18(8):605-608
- 99 Chemisch-pharmazentische Werke Bad Homburg A.-G., Bad Homburg. 1923
Herstellung eines kolloidal loslichen kieselesauregranulats
D.R. Pat. 386,760 issued Dec. 15, 1923
- 100 Chirico, M. 1946
The combined action of mechanical (kieselguhr) and chemical (3,4-benzopyrene) stimulus on connective tissue
Atti Accad. Lincei Cl. Sci. Fis., Mat. Nat. 1:239-241
- 101 Chirico, M. 1946
Hyperplasia provoked by infusorial earth in association with 3,4-benzopyrene
Tumori 20:15-19
- 102 Cipolla, Richard H., Donald W. Davis, and Carl R. Vander Linden (to Johns-Manville Corp.). 1961
Calcium silicate hydrate as an anticaking agent in dairy products
U.S. Pat. 2,995,447 issued Aug. 8, 1961
- * 103 Collet, A., and H. Moussard. 1953
Biliary elimination of silica; experimental study in the rabbit
Compt. Rend. Soc. Biol. 146:1574-1577
- * 104 Committee on Specifications. 1972
Food Chemicals Codex
Committee on Food Protection, National Research Council/National Academy of Sciences, Wash. D.C. pp. 716-717

- * 105 Comolli, R. 1965
Effect of silica on protein biosynthesis by rat liver cell-free systems
Experientia 21(9):543-544
- 106 Comolli, R. 1967
Cytotoxicity of silica and liberation of lysosomal enzymes
J. Pathol. Bact. 93:241-253
- 107 Comolli, Roberto, and Antonio Perin. 1963
The metabolism of macrophages after phagocytosis of powdered silica (Fr.)
Med. Lav. 54(5):329-334
- 108 Comrie, A.A.D. 1936
Über asche-, kiesel- und eisen-gehalt von malzen
J. Inst. Brew. 42(N.S. 33):350-353
- 109 Conning, D.M., and A.G. Heppleston. 1966
Reticuloendothelial activity and local particle disposal. A comparison of the influence of modifying agents
Brit. J. Exp. Pathol. 47(4):388-400
- 110 Coombs, J., and B.E. Volcani. 1968
Biochemistry and fine structure of silica-shell formation in diatoms. Chemical changes in the wall of *Navicula pelliculosa* during its formation
Planta 82:280-292
- 111 Coombs, J., P.J. Halicki, O. Holm Hansen, and B.E. Volcani. 1967
Studies on the biochemistry and fine structure of silica shell formation in diatoms. Changes in concentration of nucleoside triphosphates during synchronized division of *Cylindrotheca fusiformis*
Exp. Cell Res. 47(1-2):302-314
- 112 Coombs, J., P.J. Halicki, O. Holm Hansen, and B.E. Volcani. 1967
Studies on the biochemistry and fine structure of silica shell formation in diatoms. II. Changes in concentration of nucleoside triphosphates in silicon-starvation synchrony of *Navicula pelliculosa*
Exp. Cell Res. 47(1-2):315-328
- 113 Cramer, R. 1948
Talc granuloma of peritoneum
Gastroenterologia 73:129-156
- 114 Crossmon, G.C. 1964
Microscopic identification of toxic dusts
Occup. Health Rev. 16:3-7
- 115 Cruess, W.V. 1946
Experiments on lye treatments in ripe olive pickling
Fruit Prod. J. 26(3):68-70
- 116 Cullen, L.J. 1970
Seasoning mixtures
U.S. Pat. 3,505,080 issued in 1970
- 117 Cummins, S.L. 1922
Anti-bactericidal properties of colloidal silica
Brit. J. Exp. Pathol. 3:237-240
- 118 Cummins, S.L. 1922
Agglutination of washed red blood-corpuscles by colloidal silica
Brit. J. Exp. Pathol. 3:260-262
- * 119 Cummins, S.L., and C. Weatherall. 1931
The retardation of lytic processes by colloidal silica solution
Brit. J. Exp. Pathol. 12(4):239-244
- * 120 Cummins, S.L., and C. Weatherall. 1931
Effects of colloidal silica upon the growth of tubercle bacilli in blood
Brit. J. Exp. Pathol. 12(4):245-249
- 121 Cummins, S.L., and C. Weatherall. 1933
Effects of colloidal silica on experimental tuberculosis in guinea-pigs
J. Hyg. 33:295-306
- 122 Cunningham, D., and R. Hans. 1968
Food release composition
U.S. Pat. 3,397,065 issued in 1968
- 123 Cuzner, Janet. 1969
Colorimetric determination of silica in beer, wort, and brewing materials
Am. Soc. Brew. Chem., Proc. 1969:190-195
- * 124 Dale, Jean C., and E.J. King. 1953
Acute toxicity of mineral dusts
Arch. Ind. Hyg. Occup. Med. 7:478-483
- 125 Dalion, J. 1966
Apropos of a fatal case of talcoma
J. Sci. Med. Lille 84:137-144
- 126 Daniel-Moussard, H. 1954
Action in vitro of silica on the activity of alkaline phosphatase
Compt. Rend. Soc. Biol. 148:1979-1981
- 127 Daniel-Moussard, H. 1955
Action of silica on the activity of ribonuclease in vitro
Compt. Rend. Soc. Biol. 149:505-507
- 128 Daniel-Moussard, H. 1961
Action of different forms of silica on oxidative phosphorylation
Compt. Rend. Soc. Biol. 155:699-703
- * 129 Daniel-Moussard, H., and A. Collet. 1960
Influence d'agents complexants sur le développement de la fibrose expérimentale par silice chez le rat. (Influence of chelating agents on the development of experimental fibrosis by silica in the rat)
Rev. Fr. Etud. Clin. Biol. 5(3):291-293
- 130 Dany, P.-J., P. Munch, and J. Kandler. 1970
Verfahren und anlage zur gewinnung von obstsaften. (Method and device for extracting fruit juice)
W. Ger. Pat. 1,814,274 issued in 1970
- 131 Davies, J.G., A.C. Gomez, and D. Boon. 1938
Die bestimmung der löslichen kiesel-säure im rohrsafft
Int. Sugar-J. 40:105-106
- 132 De Balsac, H.F. Heim, E. Fernback, and G. Rullier. 1940
Effects, question whether silica influences development and virulence of tubercle bacillus
Bull. Acad. Med. (Paris) 123:365-368
- 133 De Lisle, August L. 1969
Storage of seeds to produce a cereal product free of frag count (insect fragments) and toxic residue
U.S. Pat. 3,476,568 issued Nov. 4, 1969
- 134 De Ritter, Elmer, L. Magid, M. Osadea, and S.H. Rubin. 1970
Effect of silica gel on stability and biological availability of ascorbic acid
J. Pharm. Sci. 59(2):229-232
- 135 Derubska, Barbara. 1967
A modification of biologic test for tuberculosis with silica-treated guinea pigs
Arch. Immunol. Ther. Exp. 15(2):204-221
- 136 Desai, Chandrakant H., Allan Burkman, and Rupert Salisbury. 1958
Chronic oral toxicity of aerated silica gel
J. Am. Pharm. Assoc. Sci. Ed. 47(8):589-590
- 137 Deutsche Gold- und Silberscheideanstalt. 1968
Fat emulsion products
Brit. Pat. 1,134,706 issued in 1968
- 138 Deutsche Gold- und Silber-Scheideanstalt vorm. Roessler. 1966
The removal of albumins from fermented beverages, such as beer
Fr. Pat. 1,424,209 issued Jan. 7, 1966

- * 139 Diffenbaugh, Willis G. 1953
Talc granulomata occurring in the peritoneal cavity. A case report
Gastroenterology 25(2):296-299
- 140 Douglass, F.M. 1947
Problem of talcum glove powder and contra-indications for its employment
Am. J. Obstet. Gynecol. 53:500-504
- 141 Dow Corning Corp. 1963.
Clarification of cane sugar juice
U.S. Pat. 3,097,970 issued Jul. 16, 1963
- 142 Dowden, Bobby F., and Harry J. Bennett. 1965
Toxicity of selected chemicals to certain animals
J. Water Pollut. Control Fed. 37(9):1308-1316
- 143 Dull, W. No date
Kieselsauretherapie, speziell
Kieselsaureinjektionen bei lungentuberkulose
Dtsch. Med. Wchschr. 49:820-821
- 144 Durocher, G. 1922
Die hydratisierte kiesel-saure. Ihre verwertung
in der chemischen industrie
Ind. Chim. 9:533-536
- 145 E.I. Du Pont De Nemours & Co. 1960
Ruminant feed compositions
U.S. Pat. 2,965,488 issued Dec. 20, 1960
- 146 Ebbrecht, E. 1931
Kieselgur in der deutschen zuckerindustrie.
Kieselgur als hilfsmittel bei entfärbungsarbeit
Dtsch. Zuckerind. 56:1191-1192, 1211-1214
- * 147 Egorova, T.S. 1967
Data on the effect of small concentrations of
silicon dioxide in the atmosphere on the health
of children (Russ.) Gyg. Sanit. 32(5):102-105
- 148 Eichhorst, H. 1920-1921
Poisoning with water-glass
Schweiz. Med. Wchschr. 50:1081; 76:275
- 149 Eiseman, B., M.G. Seelig, and N.A. Womack. 1948
Talcum powder granuloma; frequent and serious
postoperative complication
Wash. Univ. Med. Alumni Q. 12:1-12
- 150 El-Nakeeb, M.A., and R.T. Yousef. 1968
Influence of various materials on antibiotics in
liquid pharmaceutical preparations
Acta Pharm. Suec. 5(1):1-8
- 151 Emerick, R.J., L.B. Embry, and O.E. Olson. 1959
Effect of sodium silicate on the development of
urinary calculi and the excretion of various
urinary constituents in sheep
J. Anim. Sci. 18(3):1025-1030
- 152 Englebrecht, F.M. 1965
Study on the influence of cytotoxic and non-
cytotoxic substances in the genesis of pulmonary
fibrosis
S. Afr. Med. J. 39:1020-1023
- 153 Epstein, E., B. Gerstl, M. Berk, and J.P. Belber.
1955
Silica pregranuloma
Am. Med. Assoc. Arch. Dermatol. 71:645-647
- 154 Erbersdobler, Helmut, and Hermann Zucker. 1969
Stable and odorless feed concentrate containing
urea or ammonium compounds
S. Afr. Pat. 68 07,857 issued Jun. 3, 1969
- 155 Eyer, H. 1972
Water conditioning
Dtsch. Med. Wochenschr. 97:258
- 156 Fallon, J.T., and F.G. Banting. 1935
Cellular reaction to silica
Can. Med. Assoc. J. 33:404-407
- 157 Faulds, J.S. 1935
Pseudotuberculoma silicoticum
J. Pathol. Bact. 41(1):129-135
- 158 Fedorova, V.I. 1961
O deistvii pyli grafita s malym sodержaniem
dвуokisi kremniya v eksperimente.
(The experimental effect of powdered graphite
containing a small amount of silicon dioxide)
Arkhh. Patol. 23(8):62-67
- 159 Fifield, C.C. 1970
Quality characteristics of wheat treated with
four inert dusts for protection against insects
during storage
Northwest. Miller 277(12):9-14
- 160 Filippova, T.V. 1967
Level of some ash elements in sugar beet and
grape pectin (Russ.)
Fiziol.-Biokhim. Osn. Pitan. Rast. 2:164-166
- * 161 Filley, Giles F., John G. Hawley, and Geo. W.
Wright. 1945
The toxic properties of silica. I.
Bronchoconstrictor effect of colloidal silica in
isolated perfused guinea-pig lungs
J. Ind. Hyg. Toxicol. 27:37-46
- 162 Fink, Hermann, and Wilhelm Riedel. 1931
Über die adsorption von eiweisskörpern in bier an
verschiedene adsorbentien
Wchschr. Brau. 48:437-439
- 163 Fischer, W. 1964
On the demonstration of small amounts of
silicates of difficult solubility (talc)
in organic material
Zentralbl. Allg. Path. 106:394-396
- 164 Flamm, A. 1970
Talc studies
Double Liaison 17(175):157-164; (176):211-216
- * 165 Food and Agriculture Organization. 1969
Toxicological evaluation of some food colours,
emulsifiers, stabilizers, anti-caking agents
and certain other substances
Food and Agriculture Organization of the United
Nations World Health Organization pp. 9, 143-147
- 166 Food and Agriculture Organization. 1970
Current food additives legislation - Italy
Curr. Food Addit. Legis. 1970(134):9-10
- 167 Food and Agriculture Organization. 1970
Current food additives legislation - Sweden
Curr. Food Addit. Legis. 1970(132):11-14
- 168 Forman, S.A., and F. Sauer. 1962
Some changes in the urine of sheep fed a hay high
in silica
Can. J. Anim. Sci. 42(1):9-17
- 169 Forman, S.A., F. Whiting, and R. Conell. 1959
Silica urolithiasis in beef cattle. III.
Chemical and physical composition of the uroliths
Can. J. Comp. Med. Vet. Sci. 23:157-162
- 170 Fristedt, D.B. 1968
Talcosis
Nord. Hyg. Tidskr. 49:66-71
- 171 Frost, Walter S. 1950
Treatment of waters for human consumption and
iodine-diglycine hydriodide composition useful
in such treatment
U.S. Pat. 2,517,651 issued Aug. 8, 1950
- 172 Fuji Kagaku Kogyo K.K. 1969
Flow improving agent
Jap. Pat. 24,293/69 issued in 1969
- 173 Fukamauchi, Hisao, Mieko Sekiguchi, and Kazuko
Iiyoshi. 1957
Analysis using hydrofluotitanic acid-hydrogen
peroxide reagent. I. Reaction of silicic acid
with the reagent
Bunseki Kagaku 6:229-232

- 174 Furukawa, T. 1960
Effects of silica on the multiplication of animal cells in tissue culture
Jap. J. Exp. Med. 30:279-288
- 175 Gallup, Willis D., C.S. Hobbs, and H.M. Briggs. 1945
The use of silica as a reference substance in digestion trials with ruminants
J. Anim. Sci. 4(1):68-71
- * 176 Gardner, L.U., and D.E. Cummings. 1933
The reaction to fine and medium sized quartz and aluminum oxide particles, silitic cirrhosis of the liver
Am. J. Pathol. 9:751-763
- 177 Gardner, Leroy U. 1937
Silicosis und verwandte bedingungen
J. Ind. Hyg. Toxicol. 19:111-125
- 178 Gardner, Leroy U. 1938
Reaction of the living body to different types of mineral dusts with and without complicating infection
Am. Inst. Min. Met. Eng., Tech. Pub. 929:15 pp
- 179 Garino-Camina, Vittorio. 1956
(Infrared absorption of) vitreous silica
Cah. Phys. 66:29-41
- 180 Gartner, Karoly. 1960
Hungarian Kieselguhr
Brauwissenschaft 13:248-253
- 181 Gattermann, E. 1951
Problem of tissue lesions (granulomas) due to talc
Zentralbl. Chir. 76:1822-1825
- 182 Gaultier, M., Et. Fournier, and Ch. Guyotjeannin. 1961
Experimental studies on the fibrogenic effects of dust and on certain general biochemical modifications (Fr.)
Proc. Int. Congr. Occup. Health, 13th, N.Y. 1960:694-698
- 183 Gebhardt, Hans T. (to Afico S.A.). 1967
Low-calorie sweetening compositon and process
U.S. Pat. 3,320,074 issued May 16, 1967
- 184 Gedick, Peter, and Winifred Pioch. 1956
Formation of organic substances in silicon dioxide granulomas
Virchow's Arch. Pathol. Anat. Physiol. 328:513-535
- 185 Geiss, V. 1939
Experiments on the use of kieselguhr as a filter material
Wein Rebe 21:178-192
- * 186 Gel'fon, I.A., and V.I. Fedorova. 1963
Changes in the blood protein fractions and histamine under the influence of amorphous and crystalline SiO₂
Ark. Patol. 25(6):45-52
- 187 General Chemical Co. 1936
Zur reinigung von ol
Can. Pat. 358,662 issued Jun. 23, 1936
- 188 Gerbeaux, J. 1968
Aluminium and magnesium pentasilicates associated with pectin and sorbitol in gastrointestinal disorders observed in children during treatment by tuberculostatics
Actual. Hepatogastroenterol. 4:B23-B24
- 189 Gerhardt, Otto. 1937
Zur talgraffination
Seifensieder-Zeitung 64:393-395
- 190 German, W.M. 1940
Lupoid-sarcoid reaction induced by foreign body (silica)
Am. J. Clin. Pathol. 10:245-250
- 191 Gernez-Rieux, Ch., C. Voisin, and C. Aerts. 1965
Behavior of the alveolar macrophages of guinea pigs after in vitro or in vivo phagocytosis of different types of silica. Protective action of P204 (poly(vinylpyridine N-oxide)) (Fr.)
Beitr. Silikose-Forsch., Sonderb. 6:115-123
- 192 Gettler, Alexander O., and Charles J. Umberger. 1945
A colorimetric method for the determination of silica in human tissues
Am. J. Clin. Path., Tech. Sect. 9:1-3
- 193 Ghemi, F. 1963
Electrocardiographic findings in mixed pneumoconiosis caused by talc
Med. Lav. 54:817-819
- * 194 Glomme, J., C.E. Holmquist, and A. Swensson. 1958
Studies on the toxicity of silicic acid
Am. Med. Assoc. Arch. Ind. Health 17:204-209
- 195 Glomme, Jon, and Ake Swensson. 1957
Urinary excretion of silica after intraperitoneal injection in rats
Acta Med. Scand. 158:387-393
- 196 Gnadt, Otto A.P. 1946
Insulin and other antidiabetic substances and preparations
Pharmazie 1:103-107
- 197 Goerke, J. 1971
Silica gel stimulates the hydrolysis of lecithin by phospholipase A
Biochim. Biophys. Acta 248:245-253
- 198 Gol'dfel'd, M.G., E.S. Vorob'eva, and O.M. Poltorak. 1966
Comparative study of catalase activity in adsorption layers of various type
Zh. Fiz. Khim. 40(10):2594-2596
- 199 Gold, Gerald, and James A. Campbell. 1964
Einfluss von ausgewahltem U.S.P. Talcum auf die stabilitat von acetylsalicylsaeure in tabletten
J. Pharm. Sci. 53:52-54
- 200 Goldman, B.S., and F.G. Pearson. 1965
Malignant pericardial effusion. Review of hospital experience and report of a case successfully treated by talc poudrage
Can. J. Surg. 8(2):157-161
- 201 Goldstein, B., and I. Webster. 1966
Intratracheal injection into rats of size-graded silica particles
Brit. J. Ind. Med. 23:71-74
- 202 Granati, A., and G. Rombola. 1955
Toxicity, effect of hyaluronidase on experimental granuloma due to quartz
Folia Med. 38:1413-1418
- 203 Grasshoff, K. 1964
On the determination of silica in sea water.
Deep-Sea Res. 11(4):597-604
- 204 Gray, J.I., and D.G. Roberts. 1970
Retention and release of volatile food flavour compounds
J. Food Technol. 5(3):231-239
- 205 Grayson, Forrest. 1935
Prufung von reinigungsmitteln fur milchanlagen
Food Ind. 7:281-282
- * 206 Grelet, Norbert. 1939
Intraperitoneal scleroses provoked by contact with diatomaceous earth in rats deprived of vitamin A
Compt. Rend. Soc. Biol. 131:1005-1008
- 207 Grieco, F. 1936
Granulomas caused by talc following laparotomy; experimental study
Arch. Ital. Chir. 42:641-659

- 208 Grundmann, Gerhard. 1950
The effect of colloidal silicic acid powder on
the healing of wounds
Zentralbl. Chir. 75:158-169
- 209 Gye, W.E., and W.J. Purdy. 1922.
The poisonous properties of colloidal silica. I.
The effects of the parenteral administration of
large doses
Br. J. Exp. Path. 3:75-94
- 210 Gye, W.E., and W.J. Purdy. 1924
Die giftigen eigenschaften der kolloidalen
kieselsaure
Brit. J. Exp. Pathol. 5:238-250
- 211 Hadni, A., C. Bouster, E. Decamps, Jm. Munier,
and P. Poinot. 1962
Absorption spectra of some inorganic and organic
compounds in the far infrared from 50 to 1000
microns (Fr.)
Colloq. Spectros. Int., 9th, Lyons, 1961 3:11-43
- 212 Haghiri, F. 1964
Strontium-90 accumulation by some vegetable crops
Ohio J. Sci. 64:371-374
- * 213 Halonen, Pentti I., Pentti Koskelo, M.H. Frick,
and Lauri Merenmies. 1962
Metastatic carcinoid tumor associated with
complicated silicosis
Acta Med. Scand. 171:477-482
- 214 Hamada, Akira. 1964
Determination of free SiO₂ by the H₃PO₄ method. I
Ind. Health 2(3-4):181-188
- 215 Hansen, Poul M.T., and Lois Linton-Smith (to
Commonwealth Scientific and Industrial Research
Organization). 1966
Free-flowing edible fat-containing powder
U.S. Pat. 3,271,165 issued in 1966
- 216 Harington, J.S., and A.C. Allison. 1965
Lysosomal enzymes in relation to the toxicity of
silica
Med. Lav. 56(6-7):471-484
- * 217 Harley, J.D., and J. Margolis. 1961
Haemolytic activity of colloidal silica
Nature (London) 189:1010-1011
- 218 Harrand, Monique. 1957
Raman spectra of vitreous substances at high
temperature
Ann. Phys. 2(13):309-317
- 219 Hasegawa, Masaru, Sadao Hirota, and Hiroshi Ogawa.
1969
Encapsulation of oily drugs
U.S. Pat. 3,421,282 issued Jan. 14, 1969
- 220 Haude, H. 1956
As cause of foreign body granuloma in peritoneum
and endometrium
Arzneim. Forsch. 10:I/110-I/117
- 221 Haynes, D.J., and V.E. Weis. 1971
Improved frying fats
U.S. Pat. 3,619,213 issued in 1971
- 222 Heffernan, P. 1929
Silica in medicinal waters
Arch. M. Hydröl. 7:208-210
- 223 Henderson, Lowell L. 1953
Sodium salicylate in rheumatic fever; effect of
adjuvant medication
Am. J. Med. Sci. 225:480-484
- 224 Henderson, W.J. 1971
Talc and carcinoma of the ovary and cervix
J. Obstet. Gynaecol. Brit. Commonw. 78:266-272
- 225 Hendrickx, H., and H. de Moor. 1970
Flowing properties of spray dried skim-milk and
whole milk
Rijksuniv. Gent. 35(1):279-290
- 226 Heppleston, A.G., and J.A. Styles. 1967
Activity of a (rat) macrophage factor in collagen
formation by silica
Nature 214(5087):521-522
- 227 Hiyeda, Kentaro, Haruo Sakane, Yuzo Shimazaki,
Hisashi Aoki, Hiroshi Nagami, Yoshikazu
Nishimoto, and Sadao Yasutake. 1940
Hypersilikaemia (Japanese)
Soc. Path. Jap. Trans. 30:517-523
- 228 Hodel, C. 1967
Histo-enzymological findings in talc granuloma
giant cells in rats
Path. Microbiol. (Basel) 30:27-34
- 229 Hokama, Y. 1971
The reversible inhibition of catalase activity by
nucleotides and its possible relationship to
mouse liver catalase depression induced by
biological substances
Cancer Res. 31:2018-2025
- 230 Holt, P.F., and J.E.L. Bowcott. 1954
Adsorption of protein on silica surfaces
Arch. Ind. Hyg. Occup. Med. 9:503-506
- * 231 Holt, P.F., D.M. Yales, and D.H. Tomlin. 1951
Distribution of injected Si-31 in rats
Biochem. J. 48:xliv
- 232 Houghtaling, Cora I., Francis S. Houghtaling,
Nestor E. Houghtaling, and Robert W. Kilburn.
1966
Citrus fruit juice concentrates
U.S. Pat. 3,227,562 issued Jan. 4, 1966
- * 233 Hughes, Walter T., and Thomas Kalner. 1966
Massive talc aspiration. Successful treatment
with dexamethasone
Am. J. Dis. Child. 111(6):653-654
- 234 Hull, William O., Harvey Keel, John Kenney, Jr.,
and Bernard W. Gamson. 1953
Diatomaceous earth
Ind. Eng. Chem. 45:256-269
- 235 I.G. Farbenindustrie Akt.-Ges., Frankfurt. 1937
Klaren von rohen fruchtsäften, süssmosten u. dgl.
D.R. Pat. 649,164 issued Aug. 17, 1937
- 236 Inoue, Tetsuo, Toshio Hayashi, and Yoshie Saito.
1967
A view on the standard of diatomaceous earth as
food additives
Bull. Natl. Inst. Hyg. Sci. 85:148-150
- 237 International Organization for Standardization.
1968
Fruit and vegetable products. Determination of
ash insoluble in hydrochloric acid
ISO Recomm. 763-1968(E):7 pp
- 238 Ivankievicz, D., S. Takacs, and Kovacs. 1970
Orale carcinome und spurenelemente des
trinkwassers. (Oral carcinoma and
trace elements in drinking water)
Vitalst., Zivilisationskr. 15(3):97-100
- 239 James, Dinah M., and J. Marks. 1956
Mechanism of silica toxicity
J. Hyg. 54:342-350
- 240 Jaretsky, R., K. Breitwieser, and F. Neuwald.
1938
Diureseversuche mit kieselsaure und
kieselsauredrogen an ratten
Arch. Pharm. Ber. Dtsch. Pharm. Ges. 276:552-559
- 241 Jindrak, K. 1965
Talc granuloma of the endometrium
Sh. Ved. Prac. Lek. Fak. Karlov. Univ. 8:505-510
- 242 Johann Ehrenzeller, Oberburen, St. Gallen. 1930
Verfahren zum sterilisieren, altern und bleichen,
sowie zur hebung der backfahigkeit von
mahlprodukten
D.R. Pat. 512,549 issued Nov. 13, 1930

- 243 Johns-Manville Corp., New York, N.Y. 1961
Verhinderung des zusammenbackens von getrockneten
milcherzeugnissen
A. Pat. 2,995,447 issued Aug. 8, 1961
- 244 Kahane, Ernest, and Georges Antoine. 1936
Nature of the silica found in living organisms;
silica of constitution and silica of
interposition
Bull. Soc. Chim. Biol. 18:1769-1782
- * 245 Kaltenbach, J.P. 1966
Hepatic catalase changes and leucine
incorporation in talc-injected mice
Cancer Res. 26:580-584
- 246 Kaltenbach, John P., and Elizabeth B. Radeke.
1964
Comparison of tumor and talc injected mice with
respect to liver catalase
Fed. Proc. 23(2, Pt. 1):548
- 247 Kamoda, M., F. Onda, H. Ito, T. Shirasaki, T.
Miki, and T. Ando. 1968
On the formation of needle-shaped sugar crystals
Proc. Int. Soc. Sugar Cane Technol. 13:362-373
- 248 Kapral, Ales M. 1954
Beer or lager
Austral. Pat. 159,331 issued Oct. 15, 1954
- * 249 Kas'yanenko, I.V. 1964
Comparative effect of NaHCO₂ and magnesium
trisilicate on gastric secretion
Vrach. Delo 1964(9):28-31
- 250 Kazantseva, T.I. 1959
The effect of silica and silicosis on the binding
of radioactive methionine by serum
Bor'ba Silikozom, Akad. Nauk S.S.S.R., Kom.
Bor'be Silikozom, Sb. Statei 4:117-121
- 251 Kaznina, N.I. 1966
On the determination of silicon dioxide
Gig. Sanit. 31:53-54
- 252 Keil, H.L., H.H. Keil, and Victor E. Nelson. 1933
Further studies on copper and iron
Proc. Soc. Exp. Biol. Med. 30(8):1153-1155
- 253 Kellum, Gene E., and Robert C. Smith. 1967
Determination of water, silanol, and strained
siloxane on silica surfaces
Anal. Chem. 39(3):341-345
- 254 Kielhofer, E. 1948
The thermal turbidity of 1947 wines
Dtsch. Wein-Ztg. 84:293-294
- 255 King, E.J., and H. Stantial. 1933
Biochemistry of silicic acid; microdetermination
of silica
Biochem. J. 27:990-1001
- * 256 King, E.J., and T.H. Belt. 1940
Silica content of tissues
J. Pathol. Bact. 51:269-275
- * 257 King, E.J., B.D. Stacy, P.F. Holt, D.M. Yates,
and D. Pickles. 1955
The colorimetric determination of silicon in the
micro-analysis of biological material and
mineral dusts
Analyst 80:441-453
- 258 King, E.J., F. Schmidt, W. Roman, and P.R.N. Kind.
1956
The effect of silica and silicic acid on enzymes
Enzymologia 17(5/6):341-351
- 259 King, E.J., H. Stantial, and M. Dolan. 1933
Biochemistry of silicic acid; presence of silica
in tissues
Biochem. J. 27:1002-1006
- * 260 King, E.J., H. Stantial, and M. Dolan. 1933
Biochemistry of silicic acid: excretion of
administered silica
Biochem. J. 27:1007-1014
- 261 King, Earl J. 1928
The estimation of silica in tissues
J. Biol. Chem. 80(1):25-31
- * 262 King, Earl J., and Thomas H. Belt. 1938
The physiological and pathological aspects of
silica
Physiol. Rev. 18:329-365
- 263 King, Earl Judson. 1939
The biochemistry of silicic acid. VIII. The
determination of silica
Biochem. J. 33(6):944-954
- 264 King, Earl Judson, and Murray M. George. 1938
Die biochemie der kieselstaure. V. Die losung
von kieselerde und silicatstaub
in korperflussigkeiten
Biochem. J. 32:417-425
- 265 King, Earl Judson, and Murray McGeorge. 1938
Die biochemie der kieselstaure. VI. Die losung
und ausscheidung der kieselerde
Biochem. J. 32:426-433
- 266 King, Earl Judson, and Wadim Roman. 1938
Die biochemie der kieselstaure. VII. Senkung der
scheinbaren loslichkeit des siliciumdioxids in
korperflussigkeiten
Biochem. J. 32:2283-2287
- 267 Klebanov, F.G., and V.I. Bychkov. 1961
On the problem of the reaction of the peritoneum
to talc
Khirurgiia (Moscow) 37:115-118
- 268 Klimov, V.V., N.V. Kagarlitskaya, and D.P.
Shcherbov. 1960
The infrared spectrometry of inorganic
substances. IV. Absorption spectra of some
silicate minerals, 2-15 μ
Tr. Kaz. Nauchn.-Issled. Inst. Miner. Syr'ya
1960(3):312-317
- 269 Klosterkoetter, W. 1965
The action of poly(vinylpyridine N-oxide) in
vitro and in vivo (Ger.)
Beitr. Silikose-Forsch., Sonderband 6:291-298
- 270 Kment, A., and H.P. Wietek. 1957
Urinary excretion of corticoids in the guinea pig
Arzneimittel-Forschung 7:564-565
- 271 Knake, Else, and Heribert Peter. 1960
The effect of oligomeric silicic acid on tissue
cultures
Beitr. Silikose-Forsch. 68:37-49
- 272 Koburu Laboratories Co. Ltd. 1971
Granular condiments
Jap. Pat. 16,947/71 issued in 1971
- * 273 Kochmann, M., and L. Maier. 1930
The biochemistry of silicic acid. 1. Influence of
silicic acid on the body weight of growing
rats. 2. Retention of silicic acid by the rat
organism. 3. Influence of silicic acid on the
elasticity of connective tissues (Ger.)
Biochem. Z. 223(1/3):228-248
- 274 Kolbach, P., and W. Finke. 1962
Determination of the minerals of malt, wort, and
beer
Monatsschr. Brau. 15:213-215
- 275 Kolobova, K.K., and V.S. Yakovleva. 1966
Determination of Si and SiO₂ in the same material
Ogneupory 31(2):56-58
- 276 Konlechner, Heinz, and Hans Haushofer. 1955
Experiments on controlled fermentation
Mitt. Klosterneuburg Ser. A, Rebe Wein 5:263-271
- 277 Koskinen, Pentti. 1956
The neutralizing effect and antipeptic properties
of some commonly used antacids
Ann. Med. Intern. Fenn. 45(Suppl. 25):90 pp.

- 278 Krasil'shchik, V.Z. 1965
Set-up for work with a hollow cathode (Russ.)
Zavod. Lab. 31(2):251
- 279 Krauss, G. 1953
Brewing technology and economics of beer
clarification with a kieselguhr filter, mass
filter, and centrifuge
Eur. Brew. Conv., Proc. Congr. 1953:225-241
- 280 Krauss, G., and W. Sellge. 1968
Wurzelklarung mittels kieselgurfiltration.
(Clarification of wort by kieselguhr filtration)
Monatsschr. Brau. 21(8):217-220
- 281 Kraut, Heinrich. 1931
Über den kieselsäuregehalt des menschlichen
blutes und seine veränderung
durch kieselsäurezufuhr
Z. Physiol. Chem. 194:81-97
- 282 Kraut, Heinrich, and Marta Weber. 1942
Determination of silica in blood
Z. Physiol. Chem. 275:127-134
- 283 Krishnakumari, M.K., and S.K. Majumder. 1962
Modes of insecticidal action of active carbon and
clay on Tribolium castaneum
Nature 193:1310-1311
- 284 Kron, St. 1934
Zur beurteilung ungarischer schweinefette
Z. Unters. Lebensm. 68:636-642
- 285 Kung, V. 1957
Toxic effect of oil-shale dust on tissues
Izv. Akad. Nauk Est. S.S.R., Ser. Biol.
1957(1):71-73
- 286 Kvorning, Sven A. 1950
The silica content of the aortic wall in various
age groups
J. Gerontol. 5:23-25
- 287 Kysela, A. 1966
Shortened analysis of talc and steatite by
complexometric determination of calcium and
magnesium
Rev. Chim. 17(2):109
- * 288 Lagergren, C. 1962
Development of silica calculi after oral
administration of magnesium trisilicate
J. Urol. 87(6):994-996
- 289 Lalonde, E.R. 1967
Analysis of talcum granulomas
J. Oral Ther. 4:104-112
- 290 Lamina, J., and N. Kruner. 1966
The insecticidal action of highly dispersed
silica (silicic acid) on ectoparasites (Ger.)
Dtsch. Tierärztl. Wochschr. 73(6):124-129
- * 291 Langendorff, H., and K. Lang. 1967
The effect of polymeric silicic acid on renal
SiO₂ excretion in man
Z. Ernährungswiss. 8:27-32
- 292 Lanning, P.C. 1963
Silicon in rice
J. Agric. Food Chem. 11(5):435-437
- 293 Lau, A. 1959
The emission spectrum of quartz vapor
Exp. Tech. Phys. 7:104-117
- 294 Le Corvaisier, H. 1959
Composition of the nonbiological precipitates in
beer
Brasserie 148:10-12
- 295 Lecinwerk, Dr. E. Laves, Hannover. 1922
Verfahren zur herstellung kieselsäurehaltiger
kunstlicher mineralwasser und limonaden
D. R. Pat. 350,247 issued Mar. 16, 1922
- 296 Lee, C.M., Jr., and E.P. Lehman. 1947
Experiments with nonirritating glove powder (and
dangers of talc)
Surg., Gynecol. Obstet. 84:689-695
- 297 Leo, H., H. von Carnap, and W. Hesse. 1923
Über die entzündungswidrige wirkung der
kieselsäure und ihre beeinflussung durch calcium
Arch. Exp. Pathol. Pharmac. 96:133-144
- 298 Leonovich, N.V., and R.I. Guseva. 1959
The technology of processing diatomaceous
material for the filtration of beer
Tr. Vses. Nauch.-Issled. Inst. Pivovar. Prom.
1959(7):48-55
- 299 Libby, Louis H., and Lloyd F. Henderson. 1948
Process of rendering oleaginous tissue material
U.S. Pat. 2,455,374 issued Dec. 7, 1948
- 300 Lichtman, A.L., J.R. McDonald, C.F. Dixon, and
F.C. Mann. 1946
Talc granuloma
Surg., Gynecol. Obstet. 83:531-546
- 301 Lippincott, Ellis R., Alvin Van Valkenburg,
Charles E. Weir, and Elmer N. Bunting. 1958
Infrared studies on polymorphs of silica and
germania
J. Res. Natl. Bur. Stand. 61(1):61-70
- * 302 Lipworth, E., B.M. Bloomberg, and F.P. Reid. 1964
Urinary calculi containing silica: a case report
S. Afr. Med. J. 38:50-51
- 303 Little, Lawrence L. 1937
Eine einfache betriebsmethode zur bestimmung der
alkalischen bestandteile von backpulvern und
waschlösungen aus alkalimischungen
J. Dairy Sci. 20:93-99
- 304 Loblich, Hans Joachim, Werner Koch, Karl Heinz
Soffge, and Martin Wordmann. 1957
Absorption of incorporated dust. Fine-grained
dusts
Beitr. Silikose-Forsch. 50:1-10
- * 305 Loeper, Jacques, and Jacqueline Loeper. 1967
Effect of silicon compounds on experimental
atheroma
Compt. Rend. Soc. Biol. 161(3):589-591
- 306 Löffler, Johannes. 1961
Determination of small amounts of SiO₂
Glastech. Ber. 34:320-324
- 307 Lohmann, W. 1938
Die kieselsäure im wasser und mineralwasser
Dtsch. Mineralwasser-Ztg. 42:759-760
- 308 Louvrier, J., and I.A. Voinovitch. 1959
Volumetric determination of silica
Ind. Ceram. 510:243-247
- 309 Luchtrath, H. 1967
Granulomas caused by modern starchy talc for
surgical gloves
Chirurg 38:360-363
- 310 Luchtrath, H., and K.G. Schmidt. 1959
Relation of talc and steatite to asbestos and
their effects in intratracheal animal
experimentation on rats
Beitr. Silikose-Forsch. 61:60 pp
- 311 Luers, Heinrich, and Sven Jacobsen. 1933
Der einfluss kieselsäurehaltigen brauwassers auf
die hefegärung
Wschr. Brau. 50:347-348
- 312 Luna, Romulo Ochoa, and Medardo Echegaray Rosales.
1958
Methods for the determination of free silica
Ind. Miner. 17(3-4):25
- * 313 Lundgren, K.D., and A. Svensson. 1953
Experimental investigations on the significance
of the size of particles in the reaction of the
peritoneum to amorphous silica
Acta Med. Scand. 145(2):84-90

- 314 Macciotta, G. 1930
Relations between endocrine secretions and silica metabolism; cases
Clin. Ed. Ig. Infant. 5:129-143
- 15 Mackey, W.A., and J.B. Gibson. 1948
Siliceous granuloma due to talc; cause of postoperative peritoneal adhesions
Brit. Med. J. 1:1077-1078
- * 316 Macnab, Gwen, and J.S. Harington. 1967
Hemolytic activity of asbestos and other mineral dusts
Nature 214 (5087):522-523
- 317 MacGillivray, A.W., and M. Matic. 1970
Composition of South African final molasses
Proc. S. Afr. Sugar Technol. Assoc. 1970(44):81-87
- 318 MacKay, A.R., and G.A.C. Synder. 1949
Talc granuloma
Northwest Med. 48:761-763
- 319 MacQuiddy, E.L., J. Perry Tollman, W. La Towsky, and Stanley Schonberger. 1939
Tissue reaction to some carbon arch dusts
J. Ind. Hyg. Toxicol. 21(10):498-513
- 320 Maddin, C.M., and R.B. Rosene. 1960
Identification by instrumental methods of chemical compounds in water-formed deposits
Am. Soc. Test. Mater., ASTM Stand., Spec. Tech. Publ. 256:28-43
- 321 Maignon, F. No date
Über die wirkungen der elektrolyse auf die gewebsdiastasen tierischen ursprungs. Überfluss an kieselssäure in der asche
Compt. Rend. Acad. Sci. 178:654-657
- 322 Majumdar, Anil Kumar, and Siddheshwar Banerjee. 1955
Estimation of silica
Anal. Chim. Acta 13:424-426
- 323 Mancino, D. 1966
Granulomatous reactions induced by silica. I. Relationships between particles size and multinucleated giant cells
Folia Med. (Napoli) 49:85-121
- 324 Mancino, D., and R. Arienzo. 1967
Effects of pulmonary lipoprotein on the cytotoxic effects of silica. II. Inhibition of the hemolytic effects of a sub-micron amorphous silica by a rat lung surface-active lipoprotein (Ital.)
Boll. Soc. Ital. Biol. Sper. 43(23):1754-1757
- 325 Mancino, D., and R. Arienzo. 1967
Effects of pulmonary lipoprotein on the cytotoxic effects of silica. I. Inhibition of the hemolytic effect of a sub-micron amorphous silica by bovine pulmonary thromboplastic lipoprotein (Ital.)
Boll. Soc. Ital. Biol. Sper. 43(23):1752-1754
- 326 Mangiarotti, L. 1942
A new bactericide--Na metasilicate
Farm. Ital. 10:427-428
- * 327 Margolis, J. 1961
The effect of colloidal silica on blood coagulation
Austral. J. Exp. Biol. Med. Sci. 39(3):249-258
- 328 Margolis, J. 1962
Physical aspects of plasma-surface reactions
Proc. Int. Congr. Hematol., 8th, Tokyo, 1960 3:1820-1825
- * 329 Markovic, B.L., and M.D. Arambasic. 1971
Experimental chronic interstitial nephritis compared with endemic human nephropathy
J. Pathol. 103:35-40
- 330 Marks, J. 1957
The neutralization of silica toxicity in vitro
Brit. J. Ind. Med. 14:81-84
- * 331 Marks, J. 1959
Neutralization of silica toxicity in vitro
Brit. J. Ind. Med. 16:166-169
- 332 Mars, G. 1970
Behavior of silica in blood serum and in the aortic walls of subjects of various ages in health and arteriosclerosis (Ital.)
G. Gerontol. 18:645-659
- 333 Matraccia, S., D. Mancino, G. Campo, and D. Caradonna. 1962
L'influsso della temperatura ambientale sullo sviluppo delle tasche granulomatoze da silice amorfa. (The influence of the ambient temperature on the development of granulomatous sacs of amorphous silica)
Arch. "De Vecchi" Anat. Patol. Med. Clin. 38(2):309-322
- 334 Matts, S.G. Flavell, C.H.J, Swan, and J. Kelleher. 1965
Double-blind trial of bismuth aluminate and magnesium trisilicate in peptic ulceration with simultaneous gastric analysis
Brit. Med. J. 1965-I(5437):753-756
- 335 Mazurek, A.A., and L.B. Kan. 1969
Photometric determination of a silicic acid impurity in demineralized water
Zavod. Lab. 35(11):1327-1328
- * 336 McClellan, George. 1941
Rapid determination of silica in acid-insoluble silicates
Assoc. Offic. Agric. Chem. 24:111-113
- * 337 McClendon, J.P., J. Gershon-Cohen, and Henry Brody 1958
Absence of gastro-intestinal lesions in rats following ingestion of silica
Proc. Soc. Exp. Biol. Med. 97(2):330-331
- 338 McCormick, G.L., W.L. Macaulay, and G.E. Miller. 1949
Talc granulomas of eye
Am. J. Ophthalmol. 32:1252-1254
- 339 McNally, W.D., and W.L. Bergman. 1935
Silicon dioxide content of lungs of infants and of placental tissue
J. Ind. Hyg. 17:171-173
- 340 Mellin, H., U. Koch, and W. Eger. 1967
Determination of silicotic properties of dusts with different particle sizes and solubility in the liver spleen test (Ger.)
Beitr. Silikose-Forsch. 91:1-14
- 341 Merliss, R.R. 1971
Talc and asbestos contaminant of rice
J. Am. Med. Assoc. 216:2144
- * 342 Merliss, R.R. 1971
Talc-treated rice and Japanese stomach cancer
Science 173:1141-1142
- 343 Meyer, Glenn W. 1966
Amorphous diatomaceous silica in place of agar where purity of media is desired (for growing fungi)
Bull. Torrey Bot. Club 93(3):201
- 344 Micks, Don W. 1960
Susceptibility of mosquitoes to silica-gel insecticides
J. Econ. Entomol. 53(5):915-918
- 345 Migaki, G., and P.M. Garner. 1969
Talc-induced granulomas in swine
J. Am. Vet. Med. Assoc. 155(10):1595-1596

- 346 Milos, Melichar. 1936
Anwendung der sedimentationsanalyse mittels
pipette zur bestimmung der korngrosse
von pharmazeutischen produkten
Cas. Cesk. Lek. 16:73-79, 97-104
- 347 Minasian, A.M. 1970
Importance of reactivity of the organism in the
process of adhesion formation in experimental
talcosis of the abdominal cavity
Eksp. Khir. Anesteziol. 15:48-50
- * 348 Modell, W., and C. Salzman. 1941
Toxicity, pharmacology of silica; effects and
mechanism of action of fatal intravenous doses
of silicic acid
J. Lab. Clin. Med. 26:774-785
- 349 Mohn, G. 1971
Metabolism of silicic acid. 3. Resorption of
defined SiO₂ variants in rats in vivo
Beitr. Silikoseforsch. 23:227-278
- 350 Mollo, P., M. Governa, M. Dellepiane, and C.
Citati. 1965
The transfer of antigenic material by macrophages
after ingestion of crystalline silica (Ital.)
Med. Lav. 56(11):693-707
- * 351 Molnar, J.J., G. Nathenson, and S. Edberg. 1962
Fatal aspiration of talcum powder by a child
New Engl. J. Med. 266:36-37
- 352 Monori, Sandor, and Erzsebet Varga. 1962
Antimicrobial effect of some cleansing materials
used in the food industry
Budap. Musz. Egy. Elelm-kem. Tansz. Kozl.
1962:18-26
- 353 Montagna, M. 1969
Determination of silica in biological materials
by atomic absorption spectrophotometry
J. Eur. Toxicol. 2(4):207-209
- 354 Morgan, J.C., and E.J. King. 1932
Method for micro gravimetric determination of
silica in tissue
J. Biol. Chem. 95:613-620
- 355 Morris, Robert H., and Myrton F. Beeler. 1967
X-ray diffraction analysis of 464 urinary calculi
Am. J. Clin. Pathol. 48(4):413-417
- 356 Moses, Campbell. 1954
The production of renal hypertension by the
injection of finely divided silica into the
renal artery
Proc. Soc. Exp. Biol. Med. 85(4):590-592
- * 357 Mosinger, M., A. Versino, J. Jouglard-Duplay, and
J. Fondarai. 1963
Electrophoretic changes in rabbits receiving
intravenous injections of silica
Arch. Maladies Prof. Med. Trav. Secur. Soc.
24(12):881-886
- 358 Mosinger, M., H. Fiorentini, M. El. Peki, and G.
Cartouzou. 1957
Metabolism and toxicology of silica
Ann. Med. Leg. Criminol. 37:197-201
- * 359 Mosinger, M., J. Jouglard-Duplay, A. Versino, and
C. Granier. 1963
Experimental silicosis in rabbits from
intravenous administration of silica (Fr.)
Arch. Mal. Prof. Med. Trav. Secur. Soc.
24(6):503-521
- 360 Moucka, V. 1936
Talc content of rice; 2 methods for rapid
detection
Z. Unters. Lebensm. 71:175-180
- 361 Mullen, J.D. 1971
Sausage casings
U.S. Pat. 3,615,715 issued in 1971
- 362 Muller, Ernst. 1936
Der metallgehalt der gallensteine und der galle
Biochem. Z. 286:182-185
- 363 Muller, M. 1963
Studies on feeding and digestion in protozoa. VI.
The effect of ingestion of non-
nutritive particles on acid phosphatase in
Paramecium multimicronucleatum
Acta Biol. Acad. Sci. Hung. 14:209-213
- 364 Munch, R. 1971
Reaction of in-vitro cultivated cells to short-
term incubation with SiO₂ dust
Beitr. Silikoseforsch. 23:173-205
- 365 Muttelet, Vitoux and C.F. 1920
Die methode von boemer fur den nachweis von talg
im schweineschmalz
Ann. Palsif. 13:593-601
- 366 Myllarniemi, H. 1966
The effect of glove powders and their
constituents on adhesion and granuloma formation
in the abdominal cavity of the rabbit
Acta Chir. Scand. 131:312-318
- 367 Nagao, N. 1941
Experimental production of sarcoma in white rats
by combined subcutaneous administration of
kieselguhr and omicron-aminoazotoluene
Gann 35:330-332
- 368 Nagel, A.H., and T.P. Finucane. 1971
French fried products
U.S. Pat. 3,574,638 issued in 1971
- 369 Narziss, P., P. Rusitzka, and C. Pulda. 1968
Neue erkenntnisse in der bierstabilisierung. (New
findings on beer stabilization)
Brauwelt 108(100/101):1833-1843
- 370 Nash, T., A.C. Allison, and J.S. Harington. 1966
Physio-chemical properties of silica in relation
to its toxicity
Nature 210(5022):259-261
- 371 Neu, Richard. 1934
Der einfluss von kieselsauregel auf farbe und
stickstoffgehalt von sojaol
Fettchem. Umsch. 41:70-71
- * 372 Newberne, Paul M., and Robert Burton Wilson. 1970
Renal damage associated with silicon compounds in
dogs
Proc. Natl. Acad. Sci. U.S. 65(4):872-875
- 373 Niklas, H., and O. Toursel. 1941
The determination of trace elements by means of
Aspergillus niger
Bodenkd. Pflanzenernahr. 23:357-360
- 374 Nippon Volatile Oil Co. 1961
Prevention of development of rancidity of salted
fishes
Jap. Pat. 2,876('61) issued Apr. 11, 1961
- * 375 NAS/NRC. No date
Comprehensive GRAS survey report
- 376 Octave Bure, Frankreich. 1934
Entwasserung und konzentration von fruchtsaften
Fr. Pat. 759,851 issued Feb. 13, 1934
- 377 Olivier, C., I. Bertrand, and G. Carbonnet. 1951
Postoperative granuloma due to talc
Mem. Acad. Chir. 77:769-775
- 378 Ono Pharmaceutical Co. Ltd. 1971
Powdered fat product
Jap. Pat. 16,507/71 issued in 1971
- * 379 Oppenheimer, Jack H., and Walter H. Riester. 1953
Influence of cortisone on leptomeningeal reaction
induced by talc
Proc. Soc. Exp. Biol. Med. 83:844-847

- * 380 Page, R.C., R.R. Hexxner, and A. Frey. 1941
Urinary excretion of silica in humans following
oral administration of magnesium silicate
Am. J. Dig. Dis. 8(1):13-15
- 381 Pankeeva, N.E., A.S. Kostenko, and A.I.
Volkovskaya. 1969
Characteristics of compounds crystallized in the
sodium oxide-aluminum oxide-silicon dioxide-
water system at 200 and 280
Fiz.-Khim. Osn. Razlozh. Alyumosilikat.
Gidrokhim. Metod. 1969:24-37
- 382 Parkin, E.A., Enid I.C. Scott, and E.R. Varley.
1956
Insecticidal effect of Nigerian diatomite
Col. Plant Anim. Prod. (London) 5:201-207
- 383 Pater, B. No date
Arbeiten aus dem Laboratorium der
arzneipflanzenversuchsanstalt in klausenburg.
1. Bodenanalyse des versuchsfeldes
Pharm. Monatsh. 4:63-67
- * 384 Paul, J. 1960
Micro-determination of soluble silica in urine
Biochem. J. 77:202-205
- 385 Pearsall, Nancy N., and Russell S. Weiser. 1968
The macrophage in allograft immunity: I.
Effects of silica as a specific macrophage toxin
Res. J. Reticuloendothel. Soc. 5(2):107-120
- 386 Pecchiai, Luciano. 1953
Histochemical investigation on intramuscular
acidophil inclusions of cells of renal tubules
of rats orally treated by silicic anhydride
Med. Sper. 24:262-280
- 387 Pernis, B. 1963
Specificity of reaction to antigenic stimulation
in lymph nodes of immature rabbits. I.
Morphologic changes and gamma-globulin production
following stimulation with diphtheria toxoid and
silica
J. Immun. 91:541-552
- 388 Pernis, B., and M.C. Battigelli. 1955
Adsorption of serum proteins by surface of
silica; possible importance in genesis of
silicosis
Med. Lav. 46:1-13
- 389 Pernis, Benvenuto, and Fiorenzo Paronetto. 1962
Adjuvant effects of silica (tridymite) on
antibody production
Proc. Soc. Exp. Biol. Med. 110:390-392
- 390 Petcov, R., and G. Raducanu. 1957
Colorimetric determination of large amounts of
silica
Rev. Chim. (Bucharest) 8:259
- 391 Petrov, V.P. 1961
Talc as a mineral and natural resource
Tr. Inst. Geol. Rudn. Mestorozhd., Petrogr.,
Mineral. Geokhim. 1961(63):3-16
- 392 Pevtsov, G.A., and V.Z. Krasil'shchik. 1963
Analysis of impurity concentrates in a discharge
tube with a hollow cathode
Zh. Anal. Khim. 18(11):1314-1316
- 393 Piazza, R. 1931
Silica content of normal and of tuberculous lung
Riforma Med. 47:369-373
- 394 Pillai, T.N.V. 1967
Silica content in coconut (Cocos nucifera) kernel
and water
Curr. Sci. 36(24):667-668
- 395 Plews, R.W. 1970
Analytical methods used in sugar refining. (A
book)
Elsevier Publ. Co. Ltd.: Amst., Neth., Lond. UK,
and N.Y., USA 234 pp.
- 396 Pochinok, Kh. N. 1958
Determination of ash constituents in plants
Nauch. Tr. Ukr. Nauch.-Issled. Inst. Fiziol. Rast.
1958(13-14):159-171
- * 397 Policard, A., and A. Collet. 1954
Toxic and fibrosing action of submicroscopic
particles of amorphous silica
Arch. Ind. Hyg. Occup. Med. 9:389-395
- * 398 Policard, A., and A. Collet. 1954
Experimental study of renal lesions produced by
silica
J. Urol. (Paris) 60:164-171
- * 399 Policard, A., A. Collet, H. Danielmoussard, and
S. Pregermain. 1960
Deposition of silica in the mitochondria of the
urinary tube. Role of renal purification
Compt. Rend. Acad. Sci. (Paris) 251:314-315
- 400 Policard, Albert, Andre Collet, and Louis Rey.
1959
Mechanisms of the toxic action of silica. Role
of phagocytosis
Compt. Rend. 249:16-17
- 401 Pradel, E. 1970
Trial of treatment with natural purified
aluminum and magnesium silicate in diarrhea of
colectomized patients
Therapeutique 46:18-20
- 402 Prasad, J. 1965
Infrared absorption spectra of some clay minerals
and mica-type structures
Trans. Indian Ceram. Soc. 24(3):78-82
- 403 Prenant, A., J. Millot, and H. Dimitresco. 1926
Comportement du sarcome du rat en presence de
gigantome a terre d'infusoires. (Behavior of rat
sarcoma in the presence of an infusorial earth
gigantoma)
Bull. Assoc. Fr. Etud. Cancer 15(6):245-275
- 404 Prica, Milan. 1934
Über die baktericide Wirkung einiger kolloider
lösungen und ihre hygienische bedeutung
Z. Hyg. Infekt.-Kr. 116:385-396
- 405 Price, R.M. 1932
The influence of silica upon the growth of the
tubercle bacillus
Can. J. Res. 7(6):617-621
- 406 Produits Silgelac, Frankreich. 1934
Entcoffeinieren von kaffeeaufgüssen
Fr. Pat. 775,227 issued Dec. 21, 1934
- 407 Raible, K. 1968
Verfahren zur behandlung von bier zur
verhinderung des entstehens von trubungen und
zur beschleunigung seiner ausstossreife. (Method
for treating beer to prevent cloudiness and to
accelerate its clarification maturity)
W. Ger. Pat. 1,442,334 issued in 1968
- 408 Raible, K. 1970
Verfahren zur erhohung der eiweisstabilitat und
der verbesserung der biologischen haltbarkeit von
bier. (Method for increasing the protein
stability and improving the biological keeping
quality of beer)
W. Ger. Pat. 1,717,084 issued in 1970
- 409 Raible, K., and H. Bantleon. 1968
Über die filtrationseigenschaften von bier.
(Filtration of beer through kieselguhr)
Monatsschr. Brau. 21(10):277-285
- 410 Raible, K.J. 1969
Beer adsorbing agents
U.S. Pat. 3,436,225 issued in 1969
- 411 Rao, C.N.B., and V. Subrahmanyam. 1945
Adulteration of broken rice and flour with
siliceous matter
Sci. Cult. 11(2):100

- 412 Ratushnyi, G.D. 1957
The use of diatomaceous earth in wine making
Vinodel. Vinograd. S.S.S.R. 17(4):24-27
- 413 Razumovskaya, N.A. 1960
Influence of silicon compounds on the state of
cholesterol-protein complexes in animal organisms
Sb. Rabot. Silikozu, Akad. Nauk S.S.S.R., Ural.
Fil. 1960(2):203-212
- 414 Read, R.O. 1970
Keeping up with all the recent changes in Food
and Drug Regulations
Can. Food Ind. 41(3):39-41.
- 415 Rees, T.D. 1971
Eyelid deformities caused by the injection of
silicone fluid
Brit. J. Plast. Surg. 24:125-128
- 416 Reinhardt, H., P. Nauroth, and K. Achenbach. 1970
Verfahren zur Behandlung von eiweisshaltigen
flüssigkeiten mit feinverteiltem gefälltem
siliciumdioxid. (Process for treating liquids,
which contain protein, with finely divided,
precipitated SiO₂)
W. Ger. Pat. 1,517,900 issued in 1970
- 417 Rider, Joseph A. 1969
Method and composition for treatment of
gastrointestinal disorders
U.S. Pat. 3,442,189 issued Jan. 14, 1969
- 418 Rilke, Franco, and Benvenuto Pernis. 1961
Morphological and histochemical aspects of
reticuloendothelial system stimulation induced
by crystalline SiO₂ particles
Med. Lav. 52:321-330
- 419 Robock, Klaus. 1967
Luminescence measurement of different silicon
dioxide modifications for study of silicosis
origination (Ger.)
Beitr. Silikose-Forsch. 92:96 pp
- 420 Roehm, P., G. Seybold, and R. Pirtkien. 1964
Various forms of experimental stomach ulcers in
rats and the influence of drugs. I. Production
of pre-stomach ulcers in rats by glucose feeding
and drug administration
Arzneimittel-Forschung 14:47-50
- * 421 Ross, W.B., and J.M. Lubitz. 1949
Talc granuloma; survey of incidence and
significance
Ann. Surg. 130:100-112
- 422 Royston, M.G. 1969
Advances in brewing separation
Proc. Biochem. 4(9):37-38
- 423 Rudsit, K. 1926
Über leukocytaire blutveränderungen unter dem
einfluss physikalisch-chemischer prozesse
Folia Haematol. 33:95-104
- 424 Rutberg, R.A. 1953
Catalytic acceleration of the process of plasma
clotting
Dokl. Akad. Nauk S.S.S.R. 89:535-538
- 425 Rutishauser, E., and M. Bianchi. 1938
Experimentelle studie über die veränderungen der
serumphosphatase bei metallvergiftungen
Arch. Int. Pharmacodyn. Ther. 58:240-247
- 426 Sabo, I., I. Modi, and I. Sekei. 1961
Effect of silicic acid and silica on blood lipids
(Russ.)
Pat. Fiziol. Eksp. Ter. 5:74-75
- 427 Sahi, R.P. 1967
Talcum granuloma
J. Indian Med. Assoc. 48:550-551
- 428 Sajo, I. 1955
Analysis of silicates, rocks, ores, slags,
refractories, etc. I. General
Acta Chim. Acad. Sci. Hung. 6:233-241
- 429 Sakane, Haruo. 1940
Determination of silica in animal and human
tissues
J. Orient. Med. 32:95-98
- * 430 Sardi, A., and A. Tomcsanyi. 1967
Fast-neutron activation analysis of silicon in
sputum
Analyst 92:529-531
- 431 Sasaki, T., N. Nagao, and I. Otsuka. 1940
Experimental studies of sarcomas produced in
white rats by combined subcutaneous injections
of kieselguhr and omicron-aminoazotoluene
Proc. Imp. Acad., Jap. 16:493-500
- 432 Sato, Shunji. 1928
Influence of the extirpation of one kidney upon
the amyloid formation due to the intravenous
injection of sodium silicate solution in the
rabbit (Jap.)
Trans. Jap. Path. Soc. 18:240-241
- * 433 Sauer, F., D.H. Laughland, and W.M. Davidson.
1959
Silica metabolism in guinea pigs
Can. J. Biochem. Physiol 37(2):183-191
- 434 Saxen, A., and P.I. Tuovinen. 1947
Experimental and clinical observations on
granulomas caused by talc and some
other substances
Acta Chir. Scand. 96:131-151
- 435 Scheel, Lester D., E. Fleisher, and F.W. Klemperer
1953
Toxicity of silica. I. Silica solutions
Arch. Ind. Hyg. Occup. Med. 8:564-573
- 436 Schlumberger, H.G. 1952
Comparative study of reaction to injury; cellular
response to methylcholanthrene and to talc in
body cavity of cockroach (Periplaneta americana)
Am. Med. Assoc. Arch. Path. 54:98-113
- 437 Schmid, P., and W. Kleber. 1960
Absorption during clarification and filtration of
beer
Brauwissenschaft 13:238-247
- 438 Schmidt-Matthiesen, H. 1955
Silicon dioxide and reactive development of
connective tissue
Arch. Path. Anat. 327:419-459
- 439 Schmied, Otto. 1953
The filtration of beer with kieselguhr
Mitt. Versuchsstn. Garungsgewerbe 7:166-171
- 440 Schuette, H.A., and D.J. Hueninck. 1937
Mineral constituents of honey. II. Phosphorus,
calcium and magnesium
Food Res. 2:529-538
- * 441 Schulz, R.Z., and Charles R. Williams. 1942
Commercial talc--animal and mineralogical studies
J. Ind. Hyg. Toxicol. 24:75-79
- 442 Secchi, G.C. 1968
Effect of some silica dusts on erythrocyte acid
phosphatase activity
Med. Lav. 59(4):294-297
- 443 Seelig, M.G. 1948
Talcum powder evil
Am. J. Surg. 76:272-273
- 444 Setkina, O.N. 1959
Infrared spectra of minerals and their practical
application
Zap. Vses. Miner. O-va. 88:39-47
- * 445 Settle, W.R., and F. Sauer. 1960
Demonstration of siliceous deposits in the
kidneys of the guinea pig
Am. J. Vet. Res. 21:709-711

- 446 Sevc, J. 1964
Contribution to the combined effect of SiO₂ and alpha irradiation in experiments on animals
Prac. Lek. 16:268-272
- 7 Sevchenko, N.A., and V.A. Florinskaya. 1956
Reflection and transmittance spectra of the different modifications of silica in the wave length range of 7 to 24 mu
Sov. Phys. "Doklady" 1:508-511
- 448 Sevchenko, N.A., and V.A. Florinskaya. 1956
Reflection and transmittance spectra of the different modifications of silica in the wave length range of 7 to 24 mu
Dokl. Akad. Nauk S.S.S.R. 109:1115-1118
- 449 Shakhbazyan, P.A. 1962
Bactericidal action of sodium metasilicate
Tr. Yubilein. Plenuma Uch. Med. Sov., Posvyashch. 40-Letiye Ustanovleniya Sov. Vlasti Arm., Erevan. 2:115-122
- 450 Shay, Harry, S.A. Komarov, H. Siplet, and Margot Gruenstein. 1947
An evaluation of some antacid and antipeptic agents in the prevention of gastric ulceration in the rat
Am. J. Dig. Dis. 14(3):99-104
- * 451 Sheen, Robert T., and Earl J. Serfass. 1960
Recent developments in automatic colorimetric chemical analysis instruments
Ann. N.Y. Acad. Sci. 87:844-856
- * 452 Sheinfain, R. Yu., and O.P. Stas. 1969
Effect of silica xerogel storage on its adsorption properties
Zh. Prikl. Khim. (Leningrad) 42(10):2363-2365
- 453 Sidorov, T.A. 1960
Infrared spectra and the structure of some glass-forming oxides
Tr. Fiz. Inst. Im. P.N. Lebedeva, Akad. Nauk S.S.S.R. 12:225-273
- 454 Silbereisen, K., and H.F. Kersting. 1968
The use of beer stabilizers in breweries (Ger.)
Monatsschr. Brau. 21(8):221-235
- 455 Silva, R.N., and I.B. Remsen. 1970
Sugar clarification
U.S. Pat. 3,539,393 issued in 1970
- 456 Simazaki, Yuzo. 1940
Myocardial lesion caused by the continuous administration of colloidal silica
J. Orient. Med. 33:1339-1355
- 457 Simson, P.W., and A. Sutherland Strachan. 1940
A study of experimental tissue reactions following intravenous injections of silica and other dusts
Publ. S. Afr. Inst. Med. Res. 9(45):95-122
- 458 Skipin, A.I. 1958
Continuous refining of sunflower oil with sodium silicate solution
Maslob.-Zhir. Prom. 24(5):15-18
- 459 Smith, G.H. 1948
"Tuberculosis" (coexistent with carcinoma of uterus) shown to be siliceous granuloma
Brit. Med. J. 1:1078-1079
- 460 Smoot, J.L., and J.G. Willis. 1952
Benign solitary granulomatous ulceration of cecum (probably due to talc)
Va. Med. Mon. 79:454-456
- 461 Solomina, V.P. 1961
Effect of lead acetate and silica on the development of experimental skin cancer (Russ.)
Fiziol. Med. 2:55-67
- 462 Someya, Shiro, Hideo Kusawa, Hiroyuki Sakabe, and Tetsuo Ohi. 1956
Influence of various dusts on reactions of guinea pigs against BCG inoculation. I. Influence of quartz dust
Ann. Rep. Jap. Assoc. Tuberc. 1956(1):56-62
- 463 Sov, M. 1970
A classic cause, still not widely known, of postoperative peritoneal adhesions: talc
Rev. Fr. Gynecol. Obstet. 65:659-660
- 464 Spengler, O., and St. Bottger. 1931
Vergleichende versuche uber die einwirkung von verschiedenen kieselgursorten (Hyflocel, Supercel und gereinigter deutscher kieselgur) auf zuckersafte
Z. Ver. Dtsch. Zuckerind. 81:791-812
- 465 Stage, G. 1969
Wurzebehandlung aus neuer sicht. (New aspects of wort processing)
Brauereltechnik 21(18):138-143
- 466 Steenken, G. 1971
Stabilisierung mit kieselgelen und sauerstoffwerte der biere. (Use of kiesel gels for stabilization and the oxygen content of the resultant beers)
Brauwelt 111(23):455
- 467 Stone, Irwin, and Philip P. Gray. 1948
Silica and tannin in worts and beers
Am. Soc. Brew. Chem., Proc. 1948:76-94
- 468 Stone, Irwin, and Philip P. Gray. 1948
Silica and tannin in worts and beers
Wallerstein Labs. Commun. 11:301-318
- 469 Strakosch, W., and H. Wurm. 1951
Talcum endometritis following intrauterine use of globuli and still of marfanil sulfathiourate (sulfonamide mixture)
Geburtsh. Frauenheilkd. 11:1109-1114
- * 470 Strecker, F.J. 1961
Animal experiments on chemically-altered silica surfaces
Beitr. Silikose-Forsch., Sonderb. 4:121-153
- 471 Strohecker, R., R. Vaubel, and K. Breitwieser. 1935
Vorkommen und nachweis der kieselsaure in verschiedenen lebensmitteln
Z. Unters. Lebensn. 70:345-353
- 472 Stubican, V. 1957
Methods for the determination of free silica
Arh. Hig. Rada 8:166-182
- 473 Su, Goug-Jen, Nicolas F. Borelli, and Adam R. Miller. 1962
An interpretation of the infrared spectra of silicate glasses
Phys. Chem. Glasses 3(5):167-176
- 474 Sue, Pierre, Rene Michon, and Jean Merinis. 1956
Metabolism in animals of intravenously administered silica and silicates containing silicon-31
Compt. Rend. 243:1809-1811
- * 475 Sugiura, K. 1942
The relation of diet to the development of gastric lesions in the rat
Cancer Res. 2(11):770-775
- * 476 Susanna, Vittoria. 1926
Ricerche farmacologiche sull'acido silicico colloidale. (Pharmacological researches with colloidal silicic acid)
Arch. Fisiol. 24(1):1-23
- 477 Sweets Laboratories Inc. 1937
Kaugummi
A. Pat. 2,076,112 issued Apr. 6, 1937
- 478 Svensson, A., J. Glomae, and G. Bloom. 1956
Of silica particles
Am. Med. Assoc. Arch. Ind. Health 14:482-486
- 479 Swingle, A.J. 1948
Morbidity and mortality in talc granuloma; fatal case
Ann. Surg. 128:144-152

- 480 Szabo, I., and J. Mody. 1958
Experimental investigation of the dysproteinemia of silicosis
Orv. Sz. (Marosvasarhely) 6:198-203
- 1 Szabo, Istvan, J. Mody, and J. Szekely. 1961
The effect of silicic acid and silicon dioxide on blood lipids
Patol. Fiziol. Eksp. Ter. 5(3):74-75
- 482 Szymczykiwicz, Konrad, and Helena Wozniak. 1966
Effect of certain industrial dusts on the oxygen uptake capacity of the pulmonary tissue of white mice (Pol.)
Med. Pr. 17(3):185-189
- 483 Takahashi, Takeo, and Shingo Miyake. 1960
Titrimetric determination of soluble silica as molybdosilicic acid
Talanta 4:1-7
- 484 Takamura, Tsutomu, Hiromi Yoshida, and Kiyokatsu Inazuka. 1964
Infrared characteristic bands of highly dispersed silica
Kolloid-Zeitschrift 195(1):12-16
- 485 Takeda Chemical Industries, Ltd., and Yasuji Hatanura. 1965
Feed for silk worms
Fr. Pat. 1,394,330 issued Apr. 2, 1965
- 486 Tandon, S.P., H.P. Khare, and S.K. De. 1968
Talc as a growth medium for nitrite forming bacteria (Nitrosomonas)
Zentralbl. Bakt., Parasitenkd., Infektionskr. Hyg., Abt. II. 122(3):250-259
- 487 Taucins, E. 1958
Determination of mineral substances (macro- and micro-elements) in biological objects
Latv. PSR Zinat. Akad. Vestis 1958(3):49-66
- 48 Thoma, K., E. Ullmann, and E. Wolferseder. 1965
Interactions between colloidal silicic acid and drugs
Sci. Pharm., Proc., 25th 1965(2):325-330
- 489 Thoma, Karl, Elsa Ullmann, and Erich Wolferseder. 1966
Effect of adjuvants in the manufacture of pharmaceutical preparations. XII. Drug release from suspensions of colloidal silicic acid, white clay, and talc (Ger.)
Sci. Pharm. 34(3):191-202
- 490 Thompson, S.A. 1939
Development of cardio-pericardial adhesions following use of talc
Proc. Soc. Exp. Biol. Med. 40:260-261
- 491 Timmington, H. 1951
Bacterial spoilage of mixtures
Pharm. J. 167:172
- 492 Toropov, S.A., and Ts.A. Gol'dina. 1956
Apparatus for determining silicon dioxide by the thermal method
Zavod. Lab. 22:1118-1120
- * 493 Trikurakis, G.A. 1940
The goitre condition, the siliceous theory and iodine prophylaxis
Polislinico (Sez. Prat.) 47(3):87-88, 91-92, 95
- 494 Tripsa, R. 1963
Experimental studies of the action of kaolin and mixtures of kaolin and SiO₂ on the organism
J. Hyg. Epidemiol. (Praha) 7:404-414
- 495 Tronzano, L. 1966
Histopathological findings in 2 cases of pneumoconiosis due to talc
Minerva Medicoleg. 86:309-315
- * 496 Tye, M.J., K. Hashimoto, and F. Fox. 1966
Talc granulomas of the skin
J. Am. Med. Assoc. 198(13):120-122
- 497 Umezaki, Yoshimi. 1961
Colorimetric determination of silica in the presence of large quantities of phosphate. A silicomolybdate method
Nippon Kagaku Zasshi 82:847-851
- 498 Unilever Ltd. 1970
Beer haze reduction
Brit. Pat. 1,215,928 issued in 1970
- 499 Unilever N.V. 1970
Method for treating beer
Neth. Pat. 6,916,263 issued in 1970
- 500 Unilever, N. V. 1968
Beer clarification
Neth. Pat. 68 06,043 issued Oct. 29, 1968
- 501 Van Ness, John H. (to Monsanto Co.). 1966
Free-flowing fumaric and adipic acid compositions
U.S. Pat. 3,250,624 issued May 10, 1966
- 502 Varley, E.R., J. W. Du Preez, E.A. Parkin, and Enid I.C. Scot. 1956
Nigerian diatomite. Insecticidal properties for the control of stored-product insects
Col. Geol. Miner. Res. 6:176-181
- 503 Vignoli, L., B. Cristau, and R. Carboni. 1949
Au sujet de l'expertise des "tapioca". (The evaluation of tapioca)
Ann. Falsif. Fraudes 42(487/489):291-293
- 504 Villanova, R. Royo, and J. Pardo Canalis. 1933
Probable reason for immunity and predisposition to pulmonary tuberculosis; role of silicic acid content in lung: experiments with cows and sheep
Rev. Espan. Med. Cir. 16:269-271
- 505 Vratny, F., and R.B. Fischer. 1960
Effect of absorbance on Raman intensities in solids
Appl. Spectrosc. 14:76-78
- 506 Vratny, F., M. Dilling, F. Gugliotta, and C.N.R. Rao. 1961
Infrared spectra of metallic oxides, phosphates, and chromates
J. Sci. Ind. Res. (India) 20B:590-593
- 507 Vysotskii, Z.Z., V.I. Danilov, and V.V. Strelko. 1967
Properties of polysilicic acid gels and conditions for generation and development of life (Russ.)
Usp. Sovrem. Biol. 63(3):362-379
- 508 Wallace, S.A. 1947
Talc granuloma (of appendix)
Tex. State J. Med. 43:33-35
- 509 Watters, P.L. 1966
Protection of packaged food from insect infestation by the use of silica gel
J. Colloid Interface Sci. 21:94-98
- 510 Weaver, James E., and John F. Griffith. 1969
Induction of emesis by detergent ingredients and formulations
Toxicol. Appl. Pharmacol. 14(2):214-220
- * 511 Webb, J.L., R.M. Selle, and C.H. Thienes. 1942
Toxicity, absorption, accumulation and excretion of ingested silica (possible role in cataract in India)
J. Ind. Hyg. Toxicol. 24:43-45
- 512 Webber, H.F.P., and L. Taylor. 1953
Evaluation of kieselguhr as a beer filter aid
J. Inst. Brew. 59:392-397
- * 513 Webster, I., C.I. Henderson, L.W. Marasas, and D.J. Keegan. 1967
Some biologically active substances produced by the action of silica and their possible significance
Inhaled Part. Vapours II, Proc. Int. Symp., 2nd, Camb., Engl. 1965:111-119

- 514 Wee, G.C. 1966
Surgical glove powder, still a surgical hazard
Mo. Med. 63:436-439
- 515 Weinmann, W. 1941
Klarungsversuche von beerenobstmosten.
(Clarification trials with unfermented juice of small fruits)
Landwirtsch. Jahrb. 90:114
- 516 Weiss, Benjamin, and Edward A. Boettner. 1967
Commercial talc and talcosis
Arch. Environ. Health 14(2):304-308
- 517 Wendland, Gunther. 1947
Treatment of the shells of eggs with special consideration of storage at higher temperatures
Pharm. Zentralhalle 86:139-145
- 518 Werescagin, G.J. 1931
Arbeiten der standardisations-kommission der internationalen vereinigung fur theoretische und angewandte limnologie. 1, 2. Methoden der hydrochemischen analyse in der limnologischen praxis
Arch. Hydrobiol. 23(1):1-64; 23(2):167-230
- 519 Weyh, H. 1969
Ursachen, verhutung und vorausbestimmung nichthiologischer biertrubungen.
(Causes, prevention and prediction of nonbiological beer hazes.) (A review)
Braueretechniker 21(9):66-72; 21(10):74-79
- 520 Wheatley, D.N. 1966
Rapid ascites tumor production (by kieselguhr in the mouse)
Nature 209(5029):1255-1256
- * 521 Whitehouse, A.G.R. 1937
The excretion of silica in urine after oral administration of silica and silicate dusts and the solubility of silica and silicate dusts in acid and alkaline solutions at 37
J. Ind. Hyg. Toxicol. 19:590-597
- 522 Whiting, F., R. Connell, and S.A. Forman. 1958
Silica urolithiasis in beef cattle. The incidence on different rations and on range
Can. J. Comp. Med. Sci. 22:332-337
- 523 Wiech, G. 1967
X-ray spectroscopic investigation of the structure of the valence bands of silicon, silicon carbide, and silicon dioxide (Ger.)
Z. Phys. 207(5):428-445
- 524 Wijkstrom, Torgil. 1936
Methods for analysis of the mineral constituents of feeding stuffs
Medd. Centralanst. Forsoksvas, Jordbruks. No. 459
- 525 Wiland, J., J.N. Masci, and B.N. Craver. 1954
A comparison of talc and absorbable starch powders
Surg., Gynecol. Obstet. 99(3):277-280
- * 526 Williamson, Alice Polk, Russell J. Blattner, and G. Gordon Robertson. 1967
Teratogenic effects of particulate compounds in the amniotic cavity of chick embryos
Proc. Soc. Exp. Biol. Med. 124(2):524-532
- 527 Willmar, H. 1971
Erfahrungen mit modernen filtersystemen.
(Experience with modern filter systems.) (A lecture)
Brauwelt 111(51):1059-1066
- 528 Wise, B.L. 1955
Reaction of brain, spinal cord and peripheral nerves to talc and starch glove powders
Ann. Surg. 142:967-972
- 529 Wit, Hugo. No date
Die physikalische entfarbbarkeit von galle
Biochem. Z. 207:141-145
- 530 Wolff, G. 1936
Notwendigkeit einer untersuchung uber den einfluss von konservierungsmitteln in der menschlichen nahrung
Ann. Falsif. Fraudes 29:533-536
- 531 Yamamoto, Takaharu. 1928
Amyloid formation due to the injection of sodium silicate solution in rabbits (Jap.)
Trans. Jap. Path. Soc. 18:239-240
- 532 Yamene, T., K. Suzuki, T. Kaga, and Y. Takamizawa. 1968
The detrimental effects of impurities occluded in affined sugars on the sugar refining processes
Proc. Int. Soc. Sugar Cane Technol. 13:380-384
- 533 Yokoi, Yumio. 1928
Amyloid formation due to the injection of blood and sodium silicate solution in mice (Jap.)
Trans. Jap. Path. Soc. 18:237-238
- 534 Yurist, I.M., and O.I. Korotkova. 1961
Rapid methods of analysis of clay, kaolin, and talc
Zavod. Lab. 27:274-277
- * 535 Zaidi, S.H., E.J. King, C.V. Harrison, and G. Nagelschmidt. 1956
Fibrogenic activity of different forms of free silica; action of fused silica, quartz, cristobalite and tridymite on livers of mice
Am. Med. Assoc. Arch. Ind. Health 13:112-121
- 536 Zajac, J. 1966
Effect of certain chemical and mechanical factors on the formation of intraperitoneal postoperative adhesions
Rocz. Pom. Akad. Med. Swierczewski 12:371-386
- * 537 Zaks, I.F., and A. Galeniyetse. 1969
Effect of silica gel on the heparin number of blood and heparinocytes (Russ.)
Heparin 1969:68-69
- 538 Zanetti, E., and L. Piccioli. 1941
Effects, action of silica on phagocytosis
Med. Lav. 32:19
- 539 Zarzycki, J., and F. Naudin. 1961
Establishment of structural parallelisms for silica gels and binary SiO₂-X₂O glasses by infrared absorption spectroscopy (Fr.)
J. Chim. Phys. 58:830-836
- 540 Zarzycki, J., and F. Naudin. 1962
The infrared spectra of cristobalite and some isostructural compounds of SiO₂, such as GeO₂ or BeF₂; comparative study of the glass networks of the systems 2SiO₂.X₂O, 2BeF₂.XF
Proc. Int. Meet. Mol. Spectry., 4th, Bologna, 1959
3:1071-1083
- 541 Zboralski, U. 1971
Verfahren zur herstellung von kunststoffdarmen.
(Process for producing artificial casings)
W. Ger. Pat. 1,939,576 issued in 1971
- * 542 Zientara, M., and S. Moore. 1970
Fatal talc embolism in a drug addict
Human Pathol. 1(2):324-327
- 543 Zimmer, Gerhart. 1923
Silicium als reizmittel
Munch. Med. Wchschr. 70:233-236
- 544 Zisman, B., M.S. Hirsch, and A.C. Allison. 1970
Selective effects of anti-macrophage serum, silica and anti-lymphocyte serum on pathogenesis of herpes virus infection of young adult mice
J. Immunol. 104(5):1155-1159